

## **Plant disease resistance and SAR regulator protein**

### **Field of the invention**

The invention relates to broad spectrum disease resistance in plants and the  
5 identification, isolation and use of a novel regulator protein of systemic  
acquired resistance (SAR).

### **Background of the invention**

Disease resistance is a primary determinant of crop yield, and monocultures  
10 of genetically uniform plants are particularly vulnerable to attack by  
pathogens to which they have low natural resistance. A key parameter in  
plant breeding is thus the selection of plants exhibiting broad range, as well  
as specific resistance to diseases caused by infectious agents, including  
viruses, bacteria and fungi. Pathogen attack can be perceived by a host  
15 plant through the specific recognition of pathogen-derived molecules. This in  
turn elicits a rapid, localised, hypersensitive response by the plant, in the  
form of rapid necrosis at the point of pathogen attack. The host-pathogen  
interaction also induces a plant immune response known as systemic  
acquired resistance (SAR), which provides long lasting protection against a  
20 spectrum of pathogens in the uninfected parts of the plant (Yang *et al.*, 1997,  
*Genes Develop.*, 11: 1621-1639). Induction of SAR is thought to rely on the  
release of one or more signal molecules, including salicylic acid (SA), at the  
site of infection and their movement throughout the plant via the phloem.  
Perception of this systemic signal by target cells leads to the coordinate  
25 expression of a subset of pathogenesis-related (PR) genes, which contribute  
to building and maintaining disease resistance. Exogenous application of SA  
appears to be sufficient to induce SAR and PR gene expression, while  
depletion of SA, by *in planta* expression of bacterial salicylate hydroxylase  
(NahG), suppresses SAR (Gaffney *et al.*, 1993, *Science* 261: 754-756).

Genetic screens, conducted in *Arabidopsis* to select mutants in the signal transduction pathway leading to SAR, have provided a fruitful approach to identify potential positive and negative regulators of SAR. Some mutants show enhanced disease susceptibility, either due to a failure to accumulate SA, for example *eds1* (Falk *et al.*, 1999, *Proc Natl Acad Sci USA*, 96: 3292-3297), or a failure to perceive SA and induce PR gene expression, as exemplified by the *npr1* mutant (Cao *et al.*, 1997, *Plant Cell*, 88: 57-63). The *npr1* mutants (also known as the *nim1* non-inducible immunity mutant), carry mutations in a gene encoding NPR1 protein, which comprises ankyrin repeats that facilitate protein-protein interactions. NPR1 is believed to interact with basic leucine zipper transcription factors that bind and regulate expression from PR gene promoters (Zhang *et al.*, 1999, *Proc Natl Acad Sci USA*, 96: 6523-6528).

Other mutants, identified by genetic screening, display enhanced disease resistance. Lesion mimic mutants which constitutively express SAR and develop spontaneous necrotic lesions in the absence of pathogen challenge are common; however these may result from pleiotropic disruption of cellular homeostasis (Molina *et al.*, 1999, *Plant J.* 17: 667-678). Constitutive defence mutants (*cpr*) have also been found which show elevated SA levels and constitutive PR gene expression, without forming spontaneous necrotic lesions (Bowling *et al.*, 1994, *Plant Cell* 6: 1845-1857; Clarke *et al.*, 1998, *Plant Cell* 10: 557-569). PR gene expression in these *cpr* mutants is dependent on the SA signal.

Mutant screens have identified two negative regulator genes of SAR, namely *SNI1* and *MPK4*. *sni1* mutations, which cause enhanced SAR, are likely to regulate SA perception, since the *sni1* (suppressor of no-immunity) mutation can restore SAR in *npr1* mutants, which are otherwise unable to respond to SA application by inducing SAR (Dong *et al.*, 2001, *Novartis Foundation Symposium* 236: 165-173). The *Arabidopsis* *MPK4* gene encodes a

Mitogen-activated Protein kinase 4 (MPK4) that under non-pathogenic conditions, constitutively represses SAR. Mutations in the *MPK4* gene lead to increased SAR, as measured by enhanced SA levels and PR gene expression, and greater resistance to both bacterial and oomycete pathogens. (Petersen *et al.*, 2000, *Cell* 103: 1111-1120). The expression of at least 16 genes, including 8 PR genes, is significantly increased in *mpk4* mutants, consistent with a constitutive SAR phenotype, while expression of certain jasmonic acid (JA)-induced genes is blocked. The constitutive SAR of *mpk4* mutants is dependent on SA, and is abolished by *in planta* expression of bacterial salicylate hydroxylase. The *mpk4 Arabidopsis* mutant is characterised by a dwarf habit, but the plants do not form spontaneous lesions. Mutants homozygous for both *mpk4* and *npr1-1* are dwarf and constitutively express PR genes and SAR as in *mpk4* mutants, while showing the SA hypersensitivity typical of *npr1-1*, suggesting that MPK4 and NPR1 may be components of independent disease resistance pathways. Unlike NPR1, MPK4 appears to be involved in cross-talk between the JA- and SA-induced gene expression. While both MPK4 and NPR1 proteins regulate plant disease responses, they are believed to control the coordinate expression of different subsets of PR genes. Those PR genes regulated by MPK4 have been found to share similar *cis*-elements in their promoter sequences that may regulate their coordinate expression, but which are distinct from NPR1 regulated PR genes (Petersen *et al.*, 2000, *supra*). One of these elements, called a W-box, is a consensus binding-site for plant-specific WRKY transcription factors (Eulgem *et al.* 2000 *TIPS* 5: 199-206) that has been shown to act as a silencing element in the promoter of the PR1 gene (Lebel *et al.* 1998 *Plant J.* 16: 223-33)

Several approaches are proposed to enhance the broad-spectrum disease resistance of crop plants. WO 9749822 describes the isolation of the *NIM1* gene, and its expression in transgenic plants in order to increase PR gene expression and thereby enhance SAR. WO 01/66755 and WO200053762

describe the isolation of various plant homologues of the *Arabidopsis NIM1* gene and their expression in transgenic plants to enhance SAR. Similarly, WO2000028036 describes transgenic plants expressing the *NPR1* gene conferring enhanced SAR. An alternative approach to increase SAR in plants is described in WO2001002574 and involves silencing expression of the gene encoding the SNI1 negative regulator polypeptide. Silencing or blocking the activity of MPK4, a second negative regulator of SAR, in order to enhance broad resistance to plant pathogens is disclosed in WO 01/41556.

It is generally recognised that wide spread use of pesticides is a standard agricultural practise which is to the detriment of the environment, and the accumulation of their residues in ground water is a serious man-made problem. Hence there is a strong desire throughout the world to reduce agricultural dependence on chemical pesticides, and to focus on enhancing the inherent resistance of plants to disease by breeding and genetic engineering. The production of crop plants with improved broad range resistance to plant pathogens relies on the identification of plant genes and their respective proteins products, whose expression determines the level and extent of immunity to pathogen attack. In particular plant genes which are components of one of more disease resistance signalling pathway, i.e., are involved in their regulation, can provide useful tools to control the timing or level of a given defence response. The value of this approach is clearly exemplified by the examples given above, where modulated expression of SAR regulatory genes in transgenic plants can enhance resistance to various pathogens. It is preferable to modulate the expression of a positive regulator of SAR, since techniques designed up-regulate gene expression in a transgenic plant are generally more effective than those required to achieve complete silencing of gene expression. It is particularly desirable that any improvement in pathogen resistance attained in the transgenic plant is not accompanied by the formation of lesions due to a spontaneous hypersensitive response, since this will be highly disadvantageous to both

the yield and quality of the crop. It is furthermore desirable to identify genes, which can be used to increase plant resistance to a wide range of natural pathogens, without impairing the plants ability to respond to and survive other predators or environmental stresses.

5

### **Summary of the invention**

The present invention is based on the identification of a positive regulator protein of systemic acquired resistance (SAR) in plants. MKS1 is shown to be an integral component of the SAR signal transduction pathway, interacting with other components of the pathway and positively regulating SA synthesis and PR gene expression. Enhancing the expression of this plant regulator protein is shown to increase SAR in plants and to increase their resistance to pathogen attack.

Accordingly, the invention provides a transgenic plant having enhanced disease resistance and increased expression of a positive regulator of systemic acquired resistance (SAR), characterised by a transgene encoding a MAP kinase substrate 1 (MKS1) polypeptide having an amino acid sequence comprising:

20

a. MAP kinase interaction domain 1 with sequence:

IXGPRPXPLXVXXDSHXIKK and

b. transcription factor interaction domain 2 with sequence:

PVVIYXXSPKVVHXXXEFMXVQRLTG, or

25

conservatively modified variants of said domain 1 and/or domain 2 sequence, wherein X refers to any amino acid residue.

In one embodiment the transgenic plant of the invention is characterised by a transgene having a nucleic acid sequence encoding a MKS1 polypeptide comprising an amino acid sequence selected from the group: SEQ ID No. 2, 6, 10, 14, 16, 20, 26, 27, 28 and conservative variants thereof.

30

In a further embodiment the transgenic plant of the invention, is characterised by a transgene encoding a MKS1 polypeptide, said transgene comprising a nucleic acid molecule having a nucleic acid sequence selected from the group: SEQ ID No. 1, 5, 9, 13, 15, and 19.

Another embodiment of the invention is directed to the use of a nucleic acid molecule that hybridises at high stringency to a nucleic acid molecule having a nucleic acid sequence selected from the group consisting of: SEQ ID No. 1, 5, 9, 13, 15, and 19, as a transgene to produce the transgenic plant of the invention having enhanced disease resistance and increased expression of a positive regulator of systemic acquired resistance.

Furthermore the transgene of the transgenic plant of the invention may comprise a homologous promoter, or alternatively the transgene may be a chimeric gene comprising a heterologous promoter selected from the group: constitutive promoter, tissue specific promoter, and inducible promoter.

The transgenic plant of the invention includes either a dicotyledonous or a monocotyledonous plant and seed from the transgenic plant.

In a further aspect of the invention is provided a method for producing the transgenic plant of the invention, characterised by introducing an expression cassette, comprising the transgene encoding the MKS1 polypeptide, into a plant and selecting the transgenic plant and its progeny expressing said MKS1 polypeptide. Furthermore the invention encompasses a recombinant vector comprising said expression cassette and the introduction of said expression cassette into a plant through transformation or via a sexual cross with a transformed plant.

In another embodiment the invention provides a method for detecting increased expression of MKS1 polypeptide in the transgenic plant of the invention, characterised in reacting an anti-MKS1 antibody with a protein extract derived from said plant. Furthermore the invention encompasses both  
5 a polyclonal and a monoclonal anti-MKS1 antibody.

In another embodiment the transgenic plant of the invention may be used for the cultivation of a crop, wherein said crop encompasses plant biomass generated by the growth of a seed or seedling, and includes reproductive  
10 parts, e.g. seed, caryopsis, cob, or fruit and vegetative parts, e.g. leaf and tuber.

In a further embodiment the transgenic plant of the invention is used in a breeding program, wherein a plant selected in the breeding program  
15 comprises the transgene having a nucleic acid sequence encoding a MKS1 polypeptide.

#### **Brief description of the figures**

Figure 1. *Arabidopsis* MPK4 and MKS1 interacting proteins.

20 A. Yeast two hybrid screening of an *Arabidopsis* cDNA Library with MPK4 as bait (BD fusion) identified MKS1 as an interacting prey (AD fusion), and screening with MKS1 as bait identified WRKY 25 and 33 as interacting prey (AD fusion). A directed two-hybrid assay (given in *italics*) between MKS1 as bait and MPK4 as prey, confirmed their interaction. Two-hybrid assays (in  
25 *italics*) between MKS1 as bait (BD fusion) and MPK3, 5, 6 and 17 as prey (AD fusion), as well as MKS1 or MPK4 as bait (BD fusion) and WRKY26, WRKY29 or WRKY25, WRKY33 as prey (AD fusion), respectively, showed no interactions. Yeast cells in the two hybrid screen were selected on the indicated nutrient depleted growth-media (-Histidine; - Leucine; - Adenine -  
30 Tryptophan) and assayed for  $\beta$ -galactosidase ( $\beta$ -gal) reporter gene activity.

B. ClustalW alignment of the amino acid sequence of *Arabidopsis* MKS1 (Acc.No:At3g18690) and homologues or orthologues from *Brassica oleracea* (Acc.No:BoBH544707 and BoBOHBT92TR + BOGQI24TF), *Glycine max* (Acc.No:GmBE020960), *Arabidopsis* (Acc.No:At1g21326; At1g68450, 5 At2g41180, AtAL138658, At2g44340, AtT46022, At2g42140, AtAL390921) and *Oryza sativa* (Acc.No:OsCAD40925; OsBAC15955; OsAP004654, Os8360.t05160, Os8355.t00567, OsAP003260), *Nicotiana tabacum* (Ntacre169), *Zea mays* (Acc.No: ZmBM340911, ZmCC442903, ZmCC613160, ZmCC635639, ZmCC661221, ZmCC700850), *Medicago* 10 *truncatula* (Acc.No: MtAC143340.1). Identified and putative phosphorylation sites (SP) in MKS1 are indicated in italics. C-termini of the three MKS1 truncations and the Pep22 sequence are indicated above the MKS1 sequence. Aligned identical or equivalent amino acid residues are box-shaded. The consensus sequence of MKS1 is given below the alignment in 15 bold, wherein Domain1 and 2 are underlined.

Figure 2. *In vitro* interaction and phosphorylation of MKS1 by MPK4.

A. 35S methionine-labelled MPK4 (lane 1), and its binding to MKS1-GST fusion protein (lane 3), but not to GST protein alone (lane 2), following 20 separation by SDS-PAGE and detection by phosphoimager.

B. Phosphorylation assay with recombinant, full-length MKS1 (lane 1), C-terminal MKS1 truncations C1-C3, identified in Figure 1B (lanes 2-4), or positive control myelin basic protein (MBP, lane 5) and HA-tagged MPK4 immunoprecipitated from transgenic plants, analysed by SDS-PAGE and 25 phosphoimager detection. Control phosphorylation assays were performed with HA-antibody immunoprecipitates of non-transgenic, wild-type (wt) plants (lanes 6-8).

C. Phosphorylation assay with recombinant, full-length MKS1 (lane 1); mutant full-length MKS1-S30A (lane 2), MKS1 C3-truncation (lane 3), or 30 mutant MKS1-S30A C3-truncation (lane 4) and HA-tagged MPK4 immunoprecipitated from transgenic plants, and analysed as in (B).



D. Top: Phosphorylation assay with recombinant, full-length MKS1 alone (lane 1) or in the presence of increasing molar ratios of Pep22, indicated in Figure 1B (lanes 2-4) by HA-tagged MPK4, immunoprecipitated from transgenic plants, and analysed as in (B). Bottom: the phosphorylation assay (D. Top) was repeated with increasing molar ratios of a 22 amino acid peptide FLG22, as a negative control.

Figure 3. *In planta* interaction of *Arabidopsis* MKS1 and MPK4.

- 10 A. Immuno-detection of MKS1 in extracts of *E. coli* before (lane 1) and after (lane 2) IPTG induction, and in an extract of wild type *Arabidopsis* rosette leaves (lane 3) by polyclonal anti-Pep22 antibody in a Western blot (WB: pa-Pep22).
- 15 B. Immuno-detection of MKS1 immunoprecipitated (IP) with monoclonal anti-Pep22 (ma-Pep22) from wild type plant extract (lane 1) or control sample lacking plant extract (lane 2) by polyclonal antibody pa-Pep22 in a Western blot (WB: pa-Pep22).
- 20 C. Immuno-detection of HA-MPK4 by anti-HA antibody (Western blot; WB: ma-HA) in immunoprecipitates (IP) of *Arabidopsis* plant extracts using anti-Pep22 antibody, ma-Pep22 (lane 1); negative control monoclonal antibody, ma-Con (lane 2); or in a total protein plant extracts (lane 3), and a mock extract, comprising buffer and maPep22 antibody (lane 4).
- 25 D. Immuno-detection of MKS1 in a Western blot with either phosphoserine/phosphothreonine antibody ( $\alpha$ -pS/TP), or polyclonal antibody (pa-MKS1), following MKS1 immunoprecipitation from extracts of rosette leaves of wild type *Arabidopsis* (*Ler*) or *mpk4* (*mpk4*) plants using monoclonal anti-Pep22 against MKS1.

Figure 4. Transgenic *Arabidopsis* plants with modified MKS1 expression

- 30 A. Immuno-detection of MKS1 with polyclonal antibody pa-Pep22 (Western blot; WB: pa-Pep22) in extracts of 35S-MKS1 transgenic *Arabidopsis* (lane

1), wild type *Arabidopsis* Ecotype Col (wt; lane 2) and RNAi-MKS1 transgenic *Arabidopsis* (lane 3) plants.

B. Growth phenotype of wild type *Arabidopsis* Ecotype Ler (wt), 35S-MKS1 transgenic *Arabidopsis* and *mpk4* mutant *Arabidopsis* plants.

5

Figure 5. Effect of MKS1 and MPK4 on expression of defense and wounding response genes in *Arabidopsis*

A. RNA blot detection of PR1 and MKS1 mRNA in *Arabidopsis* wild type Ecotype Ler (wt; lane 1), 35S-MKS1 transgenic (lane 2) and *mpk4* mutant (lane 3) plants.

10

B. RNA blot detection of VSP and WR3 mRNA accumulation in rosette leaves from *nahGmpk4* (lanes 1-4) and wild type Ecotype Ler plants (wt; lanes 5-8), at different times after wounding.

C. RNA blot detection of VSP mRNA in rosette leaves from wild type Ecotype Col (lanes 1 and 2) and RNAi-MKS1 plants (lanes 3 and 4), at 0h and 2h after wounding.

15

D. RNA blot detection of PDF1.2 mRNA in wild type *Arabidopsis* Ecotype Ler (wt; lane 1 and 2), RNAi-MKS1 (lane 3 and 4), and 35S-MKS1 transgenic (lane 5 and 6) plants, at 0h and 48 hr after methyl jasmonate (MeJA) treatment.

20

Figure 6. Properties of *Arabidopsis* plants with altered MKS1 expression

A. Salicylate levels (ng/g FT (fresh weight)) in leaves from 4-week-old 35S-MKS1 transgenic *Arabidopsis* and wild type (wt) plants grown in soil. Error bars show standard deviation of triplicates; absence indicates insignificant differences.

25

B. Pathogen virulence assay of 4-week-old wild-type *Arabidopsis* Ecotype Ler (wt), 35S-MKS1 transgenic *Arabidopsis* and *mpk4* mutant *Arabidopsis* plants inoculated with the virulent strain DC3000 of *Pseudomonas syringae* pv. *tomato* at a concentration of  $1 \times 10^5$  colony-forming units per ml

30

(CFU/ml). Values represent average and standard deviations of cfu extracted from leaf disks in three independent samplings.

C. Pathogen virulence assay of wild type *Arabidopsis* Ecotype Col (wt) and RNAi-MKS1 transgenic *Arabidopsis* plants. Values given are as in B.

- 5 D. GFP fluorescence detection of the GFP fusion proteins: MKS1-GFP, MPK4-GFP and GUS-GFP expressed in leaf mesophyll cells of transgenic *Arabidopsis* plants using confocal microscopy. Subcellular compartments indicated are: cytoplasm (cy); nucleus (nu).
- 10 Figure 7. Suppression of *mpk4* by *MKS1*-RNAi. a. Phenotypes of wild type (*Ler*), *mpk4* carrying *MKS1*-RNAi (*mpk4*/*MKS1*-RNAi), and *mpk4*. b. RNA blot detection of *PR1* mRNA in wild type (*Ler*), *mpk4*/*MKS1*-RNAi, and *mpk4*. c. Pathogen virulence assay of 4-week-old *Arabidopsis* wild type Ecotypes *Ler* and *Col*, *mpk4*, *MKS1*-RNAi, and *mpk4*/*MKS1*-RNAi inoculated with the
- 15 virulent strain DC3000 of *Pseudomonas syringae* pv. *tomato* at a concentration of  $1 \times 10^5$  colony-forming units per ml (CFU/ml). Values represent average and standard deviations of cfu extracted from leaf disks in three independent samplings.
- 20 Figure 8. A model of defense signaling in *Arabidopsis*, highlighting MPK4, MKS1, WRKY25 and WRKY33.

## Detailed description of the invention

### I. Abbreviations

- 25 **GST:** Glutathione-S-transferase
- MKS1:** Map Kinase Substrate 1
- MPK4:** Mitogen-Activated Protein Kinase 4
- NahG:** bacterial salicylate hydroxylase
- PR gene /protein:** Pathogen Related gene/protein
- 30 **SA:** Salicylic Acid
- SAR:** Systemic Acquired Resistance

**WB:** Western Blot

**WT:** wild type

## II. Definitions

5 ***Agrobacterium*-mediated transformation:** is a technique used to obtain transformed plants by infection with *Agrobacterium tumefaciens*. During the transformation process the bacteria transfers a DNA fragment (T-DNA) from an endogenous plasmid into the plant genome. For transfer of a gene of interest the gene is first inserted into the T-DNA region of *Agrobacterium*  
10 *tumefaciens*, which is subsequently used for infection using the floral dip method according to Clough and Bent, 1998 in *Plant J* 16: 735-743.

**Antibody:** immunoglobulin protein that is produced in the body in response to immunisation with an antigen (for example MKS1 polypeptide or peptide fragment thereof), and that binds specifically to that antigen.

15 **Breeding program:** A breeding program encompasses the selection of progeny resulting from a sexual cross between parent plants. The sexual cross may be between defined parent plants or between a random population of parent plants. The progeny resulting from the cross are selected according to defined selection criteria including, but not limited to agronomic  
20 performance e.g. disease resistance, drought resistance, heat tolerance, yield, and the inheritance of a specific gene including a transgene.

**cDNA:** complementary DNA, comprising a 1<sup>st</sup> strand, complimentary to a mRNA molecule generated by reverse transcription, from which a 2<sup>nd</sup> complementary strand may be generated with a polymerase.

25 **Chimeric gene:** refers to a nucleic acid sequence, comprising a promoter operably linked to a second nucleic acid sequence containing an ORF or fused ORFs, which optionally may be operably linked to a terminator sequence. The promoter sequence is not normally operatively linked to the second nucleic acid sequence as found in nature, but is able to regulate  
30 transcription or expression of the second nucleic acid sequence. The second

nucleic acid sequence codes for a mRNA and may be expressed as a protein.

**Conservatively modified variant:** refers to a polypeptide sequence when compared to a second sequence, and includes individual conservative amino acid substitutions as well as individual deletions, or additions of amino acids. Conservative amino acid substitution tables, providing functionally similar amino acids are well known in the art. The following five groups each contain amino acids that are conservative substitutions for one another:

I: valine (V), leucine (L), isoleucine (I), methionine (M);

II: phenylalanine (F), tyrosine (Y), tryptophan (W);

III: arginine (R), lysine (K), histidine (H), glutamine (Q);

IV: aspartic acid (D), glutamic acid (E), asparagine (N), glutamine (Q);

V: alanine (A), serine (S), threonine (T).

In addition, individual substitutions, deletions or additions which alter, add or delete a single amino acid or a small percentage of amino acids in an encoded sequence are also "conservatively modified variants". When referring to nucleic acid sequences, conservative modified variants are those that encode an identical amino acid sequence, (in recognition of the fact that codon redundancy allows a large number of different sequences to encode any given protein); or conservative modified variant; or a conservative modified variant having deletions or additions of a single amino acid or a small percentage of amino acids in the encoded sequence.

**Crop:** a crop encompasses plant biomass generated by the growth of a seed or seedling, and includes reproductive parts, e.g. seed, caryopsis, cob, or fruit and vegetative parts, e.g. leaf and tuber.

**Dicotyledenous plant:** flowering plant having two cotyledons in the seed.

**Disease resistance:** the term disease resistance indicates the ability of a plant to resist pathogen attack. 'As used herein "enhanced" resistance is a greater level of resistance to a disease causing pathogen by a transgenic or genetically modified plant, produced by the method of the present invention, as compared with a non-modified, control plant. In a preferred embodiment

the level of resistance to a pathogen is at least 5%, preferably at least 10%, more preferably at least 20% greater than the resistance of a control plant.

**Exon:** protein coding sequence of a gene sequence.

**Expression cassette:** a nucleic acid sequence capable of directing

5 expression of a particular nucleotide sequence in an appropriate host cell, comprising a promoter operably linked to the nucleotide sequence of interest, which is operably linked to termination signals. The expression cassette comprising the nucleotide sequence of interest may be chimeric, meaning that at least one of its components is heterologous with respect to at least one  
10 of its other components. The expression cassette may also be one which is naturally occurring but has been obtained in a recombinant form useful for heterologous expression. Typically, however, the expression cassette is heterologous with respect to the host, i.e., the particular nucleic acid sequence of the expression cassette does not occur naturally in the host cell  
15 and must have been introduced into the host cell or its progenitor by a transformation event.

**Fusion protein:** polypeptide read-through expression product of a gene comprising two or more protein coding sequences fused in frame.

**Genetically modified plant:** in terms of the present invention relates to a  
20 non-naturally occurring plant, whose genome has been artificially modified by genetic manipulation techniques, e.g., chemical mutagenesis, site-directed mutagenesis, homologous recombination (Terada *et al.* 2002 *Nature Biotech.* 20: 1030-1034) and transformation.

**Genomic DNA:** DNA sequences comprising the genome of a cell or  
25 organism.

**Heterologous:** a polynucleotide sequence is "heterologous to" an organism or a second polynucleotide sequence if it originates from a foreign species, or from a different gene, or is modified from its original form. A heterologous promoter operably linked to a coding sequence refers to a promoter from a  
30 species, different from that from which the coding sequence was derived, or, from a gene, different from that from which the coding sequence was derived.

**Homologous:** a polynucleotide sequence is "homologous to" an organism or a second polynucleotide sequence if it originates from the same species, or gene. A homologous promoter refers to a gene promoter operably linked to the coding sequence of the same gene.

- 5    **Homologue:** is a gene or protein that is substantially identical to another gene's sequence or another protein's sequence.

**Host cell:** A prokaryotic or eukaryotic cell which may be transformed with an expression cassette cloned in a vector. The host cell may be a bacterial (for example *Agrobacterium* spp, or *E.coli*) or plant cell (for example a  
10    monocotyledenous or dicotyledenous plant cell. The protein encoded by the expression cassette may be expressed and purified from the host cell.

**Identity:** refers to nucleic acid or polypeptide sequences that are the same or have a specified percentage of nucleic acids of amino acids that are the same, when compared and aligned for maximum correspondence over a  
15    comparison window, as measured using one of the sequence comparison algorithms listed herein, or by manual alignment and visual inspection. When percentage of sequence identity is used in reference to proteins, it is recognized that residue positions that are not identical often differ by conservative amino acid substitutions, where amino acid residues are  
20    substituted for amino acid residues with similar chemical properties (e.g., charge or hydrophobicity) and therefore do not change the functional properties of the molecule. Where sequences differ in conservative substitutions, the percent sequence identity may be adjusted upwards to account for the conservative nature of the substitution. Typically this involves  
25    scoring a conservative substitution as a partial rather than a full mismatch, thus increasing the percent identity. Means for making these adjustments are well known to those skilled in the art.

**Interacting:** in terms of the present invention, relates to a physical interaction between two or more proteins, and their association for a duration sufficient  
30    to be detectable by known bioassays. For example, interacting proteins are

detected by the yeast 2-hybrid screen and assay, and by co-precipitation with antibodies with affinity to one of the interacting proteins.

**Intron:** is a non-coding sequence interrupting a protein coding sequence within a gene sequence.

- 5 **Isolated:** in the context of the present invention an isolated protein (polypeptide) or an isolated nucleic acid molecule is a protein or nucleic acid molecule that, by the hand of man, exists apart from its native environment, and is therefore not the product of nature. The isolated protein or nucleic acid molecule may exist in a purified form or in a non-native environment such as,  
10 for example, a transformed host cell.

**MAP kinase:** mitogen-activated protein kinase, which acts downstream of other MAPK kinases, in reversible phosphorylation cascades to transduce extracellular signals into cellular responses (for example MPK4, 3, 5, 6, 17).

- 15 **MKS1:** MAP Kinase Substrate1 (MKS1) polypeptide is a positive regulator of SAR and enhances plant disease resistance. The primary amino acid sequence of MKS1 comprises domain 1 with sequence:  
GPRXPXPLSVXXDSHKIKKP and domain 2 with sequence:  
PVVIYXXSPKVHXXXXEFMXVQRLTG, and conservatively modified variants thereof, wherein X refers to any amino acid residue. MKS1 is  
20 phosphorylated at one or more sites by a MAP kinase and it interacts with a transcription factor (for example a WRKY transcription factor). A MKS1 polypeptide includes a truncated or deleted fragment thereof that retains domain 1 and domain 2 sequences and the functional properties of being a positive regulator of SAR and enhancing plant disease resistance. Domain I  
25 has the functional property of comprising part or all of the interaction site for MAK kinase, while domain 2 has the functional property of comprising part or all of the interaction site for a transcription factor e.g. a WRKY-type transcription factor.

- 30 **Monocotyledenous plant:** includes, but is not limited to, barley, maize, oats, rice, rye, sorghum, wheat and members of the grass family *Poaceae*.



**Mutant:** a plant or organism with a modified genome sequence resulting in a phenotype which differs from the common wild-type phenotype.

**Native:** as found in nature, and with respect to "native promoter" refers to a promoter operably linked to its homologous coding sequence.

5    **RNA blot analysis:** a technique for the quantitative analysis of mRNA species in an RNA preparation involving size separation of RNA by agarose gel electrophoresis, subsequent transfer of RNA from the gel to a nucleic acid binding membrane, and hybridisation of the membrane with sequence specific probes.

10   **Operably linked:** refers to a functional linkage; for example between a promoter and a second sequence, wherein the promoter sequence initiates transcription of RNA corresponding to the second sequence.

**ORF:** Open Reading Frame, which defines one of three putative protein coding sequences in a DNA polynucleotide.

15   **Orthologue:** Homologous genes (or proteins) diverging concurrently with the evolutionary divergence of the organism harbouring them. Orthologues commonly serve the same function within the organisms and are most often located in a similar position on the genome.

20   **PCR:** Polymerase Chain Reaction is a technique for the amplification of a DNA polynucleotide, employing a heat-stable DNA polymerase and short oligonucleotide primers, which hybridise to the DNA polynucleotide template in a sequence specific manner and provide the primer for 5' to 3' DNA synthesis. Sequential heating and cooling cycles allow denaturation of the double-stranded DNA template and sequence-specific annealing of the  
25   primers, prior to each round of DNA synthesis. PCR is used to amplify a DNA polynucleotide employing the following standard protocol or modifications thereof:

30   PCR amplification is performed in 25 µl reactions containing: 10 mM Tris-HCl, pH 8.3 at 25°C; 50 mM KCl; 1.5 mM MgCl<sub>2</sub>; 0.01% gelatin; 0.5 unit Taq polymerase and 2.5 pmol of each primer together with template genomic DNA (50-100 ng) or cDNA. PCR cycling conditions comprise heating to 94°C

for 45 seconds, followed by 35 cycles of 94°C for 20 seconds; annealing at X°C for 20 seconds (where X is a temperature between 40 and 70°C defined by the primer annealing temperature); 72°C for 30 seconds to several minutes (depending on the expected length of the amplification product). The  
5 last cycle is followed by heating to 72°C for 2-3 minutes, and terminated by incubation at 4°C.

**Phosphorylated:** in terms of the present invention relates to the phosphorylation of a protein, such as MKS1, by a protein kinase, such as a MAP kinase. Phosphorylation sites are commonly serine and/or threonine  
10 residues on the protein. Protein kinases act to regulate the activity of proteins by covalently attaching phosphate groups. The addition of this large charged group to the protein will usually result in changes in the target protein's conformation. These conformational changes typically result in changes in the protein's activity (either up or down) or association with other  
15 proteins. Protein phosphatases act in an opposite fashion and regulate proteins by removing phosphate groups that have been covalently attached to a protein (by a protein kinase).

**Polynucleotide molecule:** or "polynucleotide", or "polynucleotide sequence" or "nucleic acid sequence" refers to deoxyribonucleotides or ribonucleotides  
20 and polymers thereof in either single- or double-stranded form. The term encompasses nucleic acids containing known analogues of natural nucleotides, which have similar binding properties as the reference nucleic acid.

**Polypeptide:** is any chain of amino acids, regardless of length or post-  
25 translational modification (for example glycosylation or phosphorylation).

**Pathogenesis Related (PR) gene:** is one that is activated or expressed in a cell of a plant in conjunction with pathogen attack and infection of the plant by a pathogen. Proteins encoded by PR genes include chitinase, extension  
(EXT1), PR1, PR5, Lipid transfer protein (LTP),  $\beta$ -1,3-glucanase  
30 (BGL2/PR2),  $\beta$ -1,3-glucanase (BGL3), glutathione-S-transferase (ERD11, PM24), LRR receptor kinase, monodehydroascorbate reductase, thionin,

osmotic, glycine-rich protein (GRR), phenylammonialyase (PAL), oxalate oxidase-like (GKP5).

**Promoter:** is an array of nucleic acid control sequences that direct transcription of an operably linked nucleic acid. As used herein, a “plant promoter” is a promoter that functions in plants. Promoters include necessary nucleic acid sequences near the start site of transcription, e.g. a TATA box element, and optionally includes distal enhancer or repressor elements, which can be located several 1000bp upstream of the transcription start site. A “tissue specific promoter” is one that specifically regulates expressed in a particular cell type or tissue, for example the promoter from the *Arabidopsis thaliana* RuBisCo small subunit gene NM\_179480 [gi:30695946]. A “constitutive” promoter is one that is active under most environmental and developmental conditions throughout the plant, for example the 35S CaMV promoter (Acc.No:V00141, J02048), the *Arabidopsis* and maize *UBI1* gene promoter (Christensen *et al.*, 1992, *Plant Mol Biol* 18: 675-689), maize *ADH* gene promoter (Last *et al.* 1991 *Theor Appl Genetics* 81: 581-588), rice *ACT1* gene promoter (McElroy *et al.* 1990 *Plant Cell* 2: 163-172). An “inducible promoter” is one which is activated in the presence of a specific agent (the inducer), which may be a chemical compound or a physical stimulus such as heat or light. The chemical compound may be one that is not found in the plant in an amount sufficient to induce activation of the inducible promoter and transcription of the operably linked gene. Examples of inducible promoters include the ecdysone agonist inducible promoter (Martinez *et al.* 1999 *Plant J.* 19: 97-106), glucocorticoid agonist inducible promoter (Aoyama and Chua, 1997 *Plant J.* 11: 605-612), copper inducible promoter (Mett *et al.* 1993 *Proc Natl Acad Sci USA* 90: 4567-4571), ethanol inducible promoter (Caddick *et al.* 1998 *Nature Biotech* 16: 177-180), tobacco *WUN1* promoter (Seibert *et al.* 1989 *Plant Cell*, 1: 961-968) and the disease-inducible *WRKY28* promoter (gi:17064157; Dong *et al.*, 2003 *Plant Mol Biol.*, 51: 21-37), and an inducible *MKS1* gene promoter may itself be used to direct expression in a *MKS1* coding sequence.

**RACE/5'RACE/3'RACE:** Rapid Amplification of cDNA Ends is a PCR-based technique for the amplification of 5' or 3' regions of selected cDNA sequences which facilitates the generation of full-length cDNAs from mRNA. The technique is performed using the following standard protocol or

- 5 modifications thereof: mRNA is reverse transcribed with RNase H<sup>-</sup> Reverse Transcriptase essentially according to the protocol of Matz *et al*, (1999) *Nucleic Acids Research* 27: 1558-60 and amplified by PCR essentially according to the protocol of Kellogg *et al* (1994) *Biotechniques* 16(6): 1134-7.

- 10 **Real-time PCR:** a PCR-based technique for the quantitative analysis of mRNA species in an RNA preparation. The formation of amplified DNA products during PCR cycling is monitored in real-time, using a specific fluorescent DNA binding-dye and measuring fluorescence emission.

- 15 **Recombinant vector:** a DNA molecule comprising sequences allowing self-replication in one or more host cells, e.g. *E.coli* or *Agrobacterium* spp., which may further comprise a heterologous chimeric gene, inserted into the vector DNA molecule. A recombinant vector, comprising a chimeric gene, may be transformed into a host cell for the purposes of expressing the chimeric gene. A recombinant vector comprising a chimeric gene also encompasses vectors for transformation of a plant, for example binary vectors.

- 20 **Regulator:** as referred to herein, is a protein which regulates another protein, pathway or response e.g. SAR, to either enhance or reduce the activity or level of said protein, pathway or response.

- 25 **SAR:** Systemic acquired resistance is a plant immune response which provides protection against a spectrum of pathogens in uninfected parts of a plant and is correlated with the expression of pathogenesis-related (PR) proteins, some with antimicrobial activity.

**Sexual cross:** refers to the pollination of one plant by another, leading to the fusion of gametes and the production of seed.

- 30 **SMART consensus:** represents the consensus sequence of a particular protein domain predicted by the Simple Modular Architecture Research Tool database (Schultz, J. *et al*. (1998)- *PNAS* 26;95(11):5857-64)

**Southern hybridisation:** A filter carrying nucleic acid (DNA or RNA) is prehybridized for 1-2 hours at 65°C with agitation in a buffer containing 7 % SDS, 0.26 M Na<sub>2</sub>HPO<sub>4</sub>, 5 % dextrane-suphate, 1 % BSA and 10µg/ml denatured salmon sperm DNA. Then a denatured, radioactively-labelled DNA probe is added to the buffer and hybridization is carried out over-night at 65°C with agitation. Unbound and non-specifically bound probe is then removed from the filter by washing. For low-stringency hybridisation, washing is carried out at 65°C with a buffer containing 2XSSC, 0.1 % SDS for 20 minutes. For medium-stringency, washing is continued at 65°C with a buffer containing 1XSSC, 0.1 % SDS for 2x 20 minutes, and for high-stringency filters are washed a further 2x 20 minutes at 65°C in a buffer containing 0.2XSSC, 0.1 % SDS. Probe labelling by random priming is performed essentially according to Feinberg and Vogelstein (1983) *Anal. Biochem.* 132(1), 6-13 and Feinberg and Vogelstein (1984) Addendum, *Anal. Biochem.*, 137(1), 266-267.

**Substantially identical:** refers to two nucleic acid or polypeptide sequences that have at least 60%, preferably 80%, most preferably 90-95% nucleotide or amino acid residue identity when aligned for maximum correspondence over a comparison window as measured using one of the sequence comparison algorithms given herein, or by manual alignment and visual inspection. This definition also refers to the complement of the test sequence with respect to its substantial identity to a reference sequence. A comparison window refers to any one of the number of contiguous positions in a sequence (being anything from between about 20 to about 600, most commonly about 100 to about 150) which may be compared to a reference sequence of the same number of contiguous positions after the two sequences are optimally aligned. Optimal alignment can be achieved using computerized implementations of alignment algorithms (e.g., GAP, BESTFIT, FASTA, and TFASTA in the Wisconsin Genetics Software Package, Genetics Computer Group, 575 Science Dr., Madison, Wis. USA) or BLAST analyses available on the site: ([www.ncbi.nlm.nih.gov](http://www.ncbi.nlm.nih.gov)). Furthermore, substantially

identical nucleic acid or polypeptide sequences perform substantially the same function.

**Transcription factor:** any protein required to initiate or regulate transcription of a gene, which may bind directly or indirectly to the DNA sequence of *cis*-elements of the gene (for example a WRKY transcription factor).

**Transgene:** refers to a polynucleotide sequence, for example a "chimeric gene", which is integrated into the genome of a plant by means other than a sexual cross, commonly referred to as transformation, to give a transgenic plant.

**Transgenic plant:** a plant harbouring a transgene stably integrated into host DNA and inherited by its progeny.

**UTR:** untranslated region of an mRNA or cDNA sequence.

**Wild type:** a plant gene, genotype, or phenotype predominating in the wild population or in the germplasm used as standard laboratory stock.

### III. Isolation of a MAP Kinase Substrate 1 protein and its homologues

The present invention concerns the protein MAP kinase substrate 1 (MKS1), isolated from *Arabidopsis thaliana*, and homologous or orthologous plant MKS1 proteins. As described more fully below in the examples, MKS1 is a positive regulator of the SAR signal transduction pathway, and plays a key role in the regulation of SA levels and PR gene expression in response to pathogen attack. MKS1 was identified by its interaction with MPK4, first detected in a yeast 2-hybrid screen. MPK4 is a negative regulator of SAR that represses SA-mediated defence responses (Petersen *et al.*, 2000, *supra*). MKS1, isolated from *Arabidopsis thaliana*, is a polypeptide of 222 amino acids residues (Seq. ID No: 2; GI:18401970; At3g18690), having 12 putative phosphorylation sites. Interaction between MPK4 and MKS1 is further demonstrated to occur *in vitro*, and *in vivo* in *Arabidopsis* plants. Interaction between MPK4 and MKS1 can furthermore lead to phosphorylation of MKS1 at one or more phosphorylation sites, where phosphorylation of residue S30 has been confirmed. MKS1, expressed as a

GFP-fusion protein, is co-localised in the nucleus of leaf mesophyll cells, together with MPK4. The targeting of MKS1, as well as MPK4, to the nucleus is consistent with its role in the SAR signal transduction pathway and induction of PR gene transcription. MKS1, isolated from *Arabidopsis thaliana*,  
5 is encoded by the intron-less gene (Seq ID No: 1; GI:18401969; At3g18690), whose function was previously unknown.

MKS1 is shown to interact with down-stream components of the SAR signal transduction pathway, which are involved in the regulation of PR gene  
10 expression. The transcription factors WRKY25 (Acc.No:GI:15991726) and WRKY33 (Acc.No:GI:21105639) are identified as interaction partners of MKS1 by 2-hybrid screening and directed 2-hybrid assay. These transcription factors are Group 1 members of a large family of WRKY plant transcription factors, which are characterised by a N-terminal WRKY domain having the  
15 conserved amino acid sequence WRKYGQK, together with a zinc finger motif (Eulgem *et al.* 2000, *Trends in Plant Sci* 5: 199-206). WRKY proteins bind to highly conserved *cis*-acting W box elements (T)(T)TGAC(C/T), which are present in defence response genes, including PR1. Although the evidence for a role of WRKY transcription factors in regulating plant defence responses is  
20 convincing, the function of the majority of members of the WRKY family is yet to be elucidated. The phosphorylation of MKS1 by MPK4, combined with the protein-protein interaction between MPK4 and MKS1 and between MKS1 and WRKY25 and 33, clearly establish MKS1 as a key regulatory protein in the SAR signal transduction pathway.

25 SAR is a plant defence mechanism, which is widely conserved in the plant kingdom (Durner J. *et al.*, 1997 *Trends in Plant Science* 2: 266-274). Thus MKS1 homologues, which function as regulator proteins in the SAR signal transduction pathway, may be found in other plants, including crop plants.  
30 MKS1 homologues and orthologues can be identified by a standard protein-protein BLAST or tblastn search against the database www:

ncbi.nlm.nih.gov/blast/BLAST.cgi . Since the isolated *Arabidopsis* MKS1 is encoded by the *MKS1* gene sequence At3g18690 (GI:18401969), an nblastn search may similarly be performed to identify plant genes encoding MKS1 homologues and orthologues. The application of this approach is illustrated in

5 the Examples, where MKS1 homologues or orthologues are identified in *Arabidopsis* (Seq ID No: 6; At1g21326), *Brassica oleracea* (Seq ID No: 10 and 14), *Glycine max* (Seq ID No: 16), and *Oryza sativa* (rice) (Seq ID No: 20), encoded by *MKS1* gene homologues or orthologues in *Arabidopsis* (Seq.ID.No:5; GI:22329704), *Brassica oleracea* (Seq ID No: 9, GI:17796488,

10 BoBH544707; Seq ID No: 13, BoBOHBT92TR + BOGQI24TF), *Glycine max* (Seq ID No: 15; GI:8283399, GmBE020960 ), and *Oryza sativa* (rice) (Seq ID.No:19, OsCAD40925;), respectively. Additional MKS1 homologues or orthologues are found in rice (Seq ID No: 26, OsAP004654) maize (*Zea mays*) (Seq ID No: 27, ZmCC613160; Seq ID No: 28, ZmCC635639),

15 tobacco (*Nicotiana tabacum*) and clover (*Medicago truncatula*) as exemplified in figure 1B. In an alternative approach, nucleotide sequences encoding plant MKS1 homologues or orthologues can be identified in libraries constructed from plant genomic or cDNA by hybridisation screening with a polynucleotide probe comprising 20 or more consecutive nucleotides of an

20 *MKS1* gene (for example At3g18690). Hybridisation screening is performed according to standard protocols, under conditions defined above. Plant genomic or cDNA may also be screened for nucleotide sequences encoding plant MKS1 homologues or orthologues by PCR, using primer sequences comprising 15 or more consecutive nucleotides of an *MKS1* gene (for

25 example At3g18690), and a standard PCR amplification protocol as defined above. The PCR amplification of nucleotide sequences encoding MKS1 can also be performed using degenerate primers whose design is based on conserved amino acid sequences in MKS1, which can be identified by ClustalW alignment of MKS1 homologues or orthologues, as shown in the

30 Examples. In the case that a MKS1 cDNA sequence is a partial sequence,



the corresponding full-length MKS1 cDNA may be generated using 5' and 3' RACE as defined above.

5 A MKS1 protein homologue or orthologue is characterised by a primary sequence that comprises domain 1 with sequence:

IXGPRPXPLXVXXDSHXIKK and domain 2 with sequence:

PVVIYXXSPKVHXXXXEFMXVQRLTG, [wherein X refers to any amino acid residue], or conservatively modified variants of said domain sequences.

10 Domain 1 of MKS1 has the functional property of interacting with the MAP kinase (e.g. MPK4) and comprising a serine-proline phosphorylation site that is phosphorylated (e.g. by MPK4), consistent with its sequence homology to a MAPK docking site (Sharrocks *et al.*, 2000, Trends Biochem. Sci. 25: 448-453). Domain 2 has the functional property of interacting with a transcription factor, in particular a WRKY-type transcription factor.

15 A MKS1 protein homologue or orthologue is substantially identical to a MKS1 protein with Seq ID No: 2, 6, 10, 14, 16, 20, 26, 27 or 28, furthermore comprising amino acid sequence domains 1 and 2, (given above) or conservatively modified variants thereof. A nucleic acid molecule encoding a MKS1 protein homologue or orthologue is characterised by a nucleotide  
20 sequence that is a substantially identical to a nucleic acid molecule with Seq ID No: 1, 5, 9, 13, 15 or 19, or more preferably a conservatively modified variant thereof. Furthermore, a MKS1 protein homologue or orthologue is characterised by the properties of being a positive regulator of SAR, enhancing plant disease resistance, being phosphorylated by a MAP kinase  
25 and interacting with a transcription factor regulating SAR gene expression, e.g. WRKY transcription factor. Phosphorylation of MKS1 by a MAP kinase can be detected by *in vitro* phosphorylation assay as illustrated in the Examples. Interaction of MKS1 with a transcription factor can be detected by yeast 2-hybrid screens and directed 2-hybrid assays as illustrated in the  
30 Examples.

#### IV Transgenic plants with modified expression of MKS1 protein

A nucleic acid molecule encoding MKS1 protein can be used to modify and enhance MKS1 protein expression in a transgenic plant of the invention and thereby induce a SAR response and increase the pathogen resistance in the plant. The MKS1 encoding nucleic acid molecule and the method provided by the invention can be utilised to induce SAR and confer disease resistance in a wide variety of plants. These plants include a monocotyledenous crop plant such as barley, maize, oats, rice, rye, sorghum and wheat; and a member of the grass family of Poaceae, such as *Phleum* spp., *Dactylis* spp., *Lolium* spp., *Festulolium* spp., *Festuca* spp., *Poa* spp., *Bromus* spp., *Agrostis* spp., *Arrhenatherum* spp., *Phalaris* spp., and *Trisetum* spp., for example, *Phleum pratense*, *Phleum bertolonii*, *Dactylis glomerata*, *Lolium perenne*, *Lolium multiflorum*, *Lolium multiflorum westervoldicum*, *Festulolium braunii*, *Festulolium loliaceum*, *Festulolium holmbergii*, *Festulolium pabulare*, *Festuca pratensis*, *Festuca rubra*, *Festuca rubra rubra*, *Festuca rubra commutata*, *Festuca rubra trichophylla*, *Festuca duriuscula*, *Festuca ovina*, *Festuca arundinacea*, *Poa trivialis*, *Poa pratensis*, *Poa palustris*, *Bromus catharticus*, *Bromus sitchensis*, *Bromus inermis*, *Deschampsia caespitosa*, *Agrostis capilaris*, *Agrostis stolonifera*, *Arrhenatherum elatius*, *Phalaris arundinacea*, and *Trisetum flavescens*.; and a dicotyledenous plant, such as alfalfa, carrot, cotton, potato, sweet potato, oilseed rape, radish, soybean, sugarbeet, sugar cane, sunflower, tobacco, turnip; vegetables such as asparagus, bean, carrot, chicory coffee, celery, cucumber, eggplant, fennel, leek, lettuce, garlic, onion, papaya, pea, pepper, spinach, squash, pumpkin, tomato; vegetable brassicas such as brussel sprouts, broccoli, cabbage, cauliflower; fruits, such as avocado, banana, blackberry, blueberry, grapes, mango, melon, nectarine, orange, papaya, pineapple, raspberry, strawberry; rosaceous fruits such as apple, apricot, peach, pear, cherry, plum and quince; herbs such as anise, basil, bay laurel, caper, caraway, cayenne pepper, celery, chervil, chives, coriander, dill, horseradish, lemon balm, liquorice, marjoram, mint, oregano, parsley, rosemary, sesame, tarragon and thyme; woody species,

such as eucalyptus, oak, pine and poplar. The coding sequence of an *MKS1* gene can be amplified by PCR using sequence specific primers, for example: *Arabidopsis MKS1* (Seq ID No: 1) is amplified by the primer pair (Seq ID No: 3 & 4); *Arabidopsis MKS1* (Seq ID No: 5) is amplified by the primer pair (Seq ID No: 7 & 8); *Brassica oleracea MKS1* (Seq ID No: 9) is amplified by the primer pair (Seq ID No: 11 & 12); *Oryza MKS1* (Seq ID No: 19) is amplified by primer pair (Seq ID No: 21 & 22). *Glycine max MKS1* (Seq ID No: 15) comprises coding sequence for a part of MKS1 protein, and the complete MKS1 coding sequence may be generated by 5' and 3'RACE, as described above using primers for 3' extension (Seq ID.No: 17) and 5' extension (Seq ID No: 18). An expression cassette is constructed comprising a nucleic acid sequence encoding a MKS1 polypeptide, substantially identical to protein SEQ ID No: 2, 6, 10, 14, 16 or 20 and furthermore comprising a domain 1 (IXGPRPXPLXVXXDSHXIKK) and domain 2 (PVVIYXXSPKVHXXXXEFMXVQRLTG) amino acid sequence or conservatively modified variants thereof, wherein said nucleic acid sequence is operably linked to a heterologous or homologous promoter and 3' terminator. The expression cassette can be transformed into a selected host plant using a number of known methods for plant transformation. By way of example, the expression cassette can be cloned between the T-DNA borders of a binary vector, and integrated into an *Agrobacterium tumerfaciens* host by transformation, and used to infect and transform a host plant (Hinchee *et al* 1988 *Bio/Technol.* 6:915-922, Ishida *et al.*, 1996 *Nat Biotechnol.* 14:745-50). The expression cassette is commonly integrated into the host plant in parallel with a selectable marker gene giving resistance to an herbicide or antibiotic, in order to select transformed plant tissue. Stable integration of the expression cassette into the host plant genome is mediated by the virulence functions of the *Agrobacterium* host. Binary vectors and *Agrobacterium tumefaciens*-based methods for the stable integration of expression cassettes into the majority of all dicotyledenous and monocotylenous crop plants are known, as described for example for rice (Hiei *et al.*, 1994, *The*

*Plant J.* 6: 271-282) and maize (Yuji *et al.*, 1996, *Nature Biotechnology*, 14: 745-750). Alternative transformation methods, based on direct transfer can also be employed to stably integrate expression cassettes into the genome of a host plant, as described by Miki *et al.*, 1993, "Procedure for introducing foreign DNA into plants", In: *Methods in Plant Molecular Biology and Biotechnology*, Glick and Thompson, eds., CRC Press, Inc., Boca Raton, pp 67-88). Promoters to be used in the expression cassette of the invention include constitutive promoters, for example the 35S CaMV promoter (Acc.No:V00141, J02048), the *Arabidopsis* and maize *UBI1* gene promoter (Christensen *et al.*, 1992, *Plant Mol Biol* 18: 675-689), maize *ADH* gene promoter (Last *et al.* 1991 *Theor Appl Genetics* 81: 581-588), rice *ACT1* gene promoter (McElroy *et al.* 1990 *Plant Cell* 2: 163-172). In an alternative embodiment, an inducible promoter may be used in the expression cassette to direct *MKS1* expression. Examples of suitable inducible promoters include the ecdysone agonist inducible promoter (Martinez *et al.* 1999 *Plant J.* 19: 97-106), glucocorticoid agonist inducible promoter (Aoyama and Chua, 1997 *Plant J.* 11: 605-612), copper inducible promoter (Mett *et al.* 1993 *Proc Natl Acad Sci USA* 90: 4567-4571), ethanol inducible promoter (Caddick *et al.* 1998 *Nature Biotech* 16: 177-180), tobacco *WUN1* promoter (Seibert *et al.* 1989 *Plant Cell*, 1: 961-968) and the disease-inducible *WRKY28* promoter (gi:17064157; Dong *et al.*, 2003 *Plant Mol Biol.*, 51: 21-37). Additionally, the inducible *MKS1* gene promoter may itself be used to direct expression in the *MKS1* expression cassette. An example of a suitable tissue-specific promoter includes the promoter from the *Arabidopsis thaliana* RuBisCo small subunit gene NM\_179480 [gi:30695946]. A terminator that may be used in the expression construct can for instance be the NOS terminator (Acc No: NC\_003065) (SEQ ID No: 24), the terminator of the *Arabidopsis thaliana* RuBisCo small subunit gene NM\_179480 [gi:30695946]. The recombinant vector comprising the *MKS1* expression cassette is optionally transformed into a plant cell together with a selectable marker gene which is located on the same or a separate recombinant vector. Marker genes that facilitate

selection of transformed plant cells, may encode peptides providing resistance to herbicide, antibiotic or drug resistance, for example, resistance to protoporphyrinogen oxidase inhibitors, hygromycin, kanamycin, G418, gentamycin, lincomycin, methotrexate, glyphosate, phosphinothricin.

- 5    Optionally, host plants transformed with an expression cassette encoding a MSK1 protein, can be crossed with a second non-transgenic plant and progeny expressing said MKS1 protein can then be selected and used in the invention.
- 10   Transgenic plants comprising a transgene expressing a MKS1 polypeptide can be used in a breeding program, in order select plants with enhanced agricultural performance that have inherited the transgene. Transgenic plants as well as plant progeny selected in such a breeding program may be
- 15   vegetative plant parts, e.g. leaf or tuber, or reproductive parts including seed, caryopsis, cob or fruit.

#### **V Plant disease resistance of transgenic plants with modified MKS1 expression**

- 20   The expression of MKS1 in transgenic plants, transformed with a MKS1 expression cassette, will be determined by the promoter to which the MKS1 coding sequence is operably linked. Where MKS1 expression is placed under the control of a constitutive promoter, MKS1 will be expressed throughout the plant at all developmental stages. The expression pattern of MKS1 will in turn
- 25   determine the SAR response pattern in the plant and the level of resistance to plant pathogen attack. Since MKS1 induces SA synthesis, all basal pathogen resistance mechanisms induced by SA will be up-regulated by MKS1 expression. Since MKS1 does not regulate expression of jasmonate-induced genes, its expression in transgenic plants will not impair jasmonate-
- 30   dependent wound responses in a plant. Furthermore, since MKS1 appears to act upstream of NPR1 in the SAR signal transduction pathway, it is expected

- to regulate a broader range of disease responses in a plant. Methods for assessing plant pathogen resistance are well known (Jach *et al.* 1995 *Plant J.* 8: 97-109 ; Whalen *et al.* 1991 *Plant Cell* 3: 49-60), and may be adapted according to the principal pathogens of the transgenic plant species. One
- 5 method for assessing the resistance of a transgenic *Arabidopsis* plant, transformed with a MKS1 expression cassette, to a bacterial pathogen (*Pseudomonas syringae*) attack is given in the Examples. Other methods for evaluating disease resistance in plants are described by Crute *et al* 1994, *Arabidopsis*, Cold Spring Harbor Press, pp 705-747. Other examples of plant
- 10 pathogens include the bacterial pathogens, *Erwinia* (for example *E. carotovora*), *Xanthomonas* (for example *X. campestris* and *X. oryzae*). Examples of fungal or fungal-like disease causing pathogens include *Alternaria*, *Ascochyta*, *Botrytis*, *Cercospora*, *Colletotrichum*, *Diplodia*, *Erysiphe*, *Fusarium*, *Gaeumanomyces*, *Helminthosporium*, *Macrophomina*,
- 15 *Nectria*, *Perenospora parasitica*, *Phoma*, *Phymatotrichum*, *Phytophthora*, *Plasmopara*, *Podosphaera*, *Puccinia*, *Puthium*, *Pyrenophora*, *Pyricularia*, *Pythium*, *Rhizoctonia*, *Scerotium*, *Sclerotinia*, *Septoria*, *Thielaviopsis*, *Uncinula*, *Venturia* and *Verticillium*.
- 20 The level of SAR in the transgenic plant can also be assessed by measuring the level of SA in the transgenic plant leaves, and the level of PR gene induction. Steady-state levels of PR mRNA can be quantitated by RNA blot hybridisation or alternatively by real-time PCR, as defined above. Application of these methods to the detection and quantitation of SAR in transgenic
- 25 plants expressing MKS1 constitutively is illustrated in the Examples.

## VI Isolated MKS1 and specific MKS1 antibodies

- A nucleic acid molecule encoding the MKS1 protein can be operably linked to a promoter sequence to form a chimeric gene capable of directing expression
- 30 of the MKS1 protein in a host cell. The nucleic acid molecule encoding MKS1 protein (ORF) can be fused in frame with a nucleic acid sequence encoding a

tag. The expression of MKS1 as a fusion protein comprising a tag (e.g. 6x histidine tag, or a glutathione-S-transferase tag) facilitates the purification of the expressed MKS1 protein. Affinity purification of tagged protein is well known in the art, and its application to the purification of MKS1 protein is described in the Examples. The chimeric gene can be cloned, as an expression cassette, in a recombinant vector, and transformed into a host cell. The expression cassette can be transformed into a bacterial cell e.g. *E.coli* and expression of tagged MKS1 protein can be controlled by an inducible promoter system, e.g. IPTG inducible promoters. Alternatively, an expression cassette can be transformed into a host plant cell, and transformed plants comprising the expression cassette can be selected. Protein extracts, prepared from tissue of the transformed plant expressing tagged-MKS1 protein, can be used for the affinity purification of tagged-MKS1.

Tagged-MKS1, MKS1, or peptide fragments thereof, can be used for the production of specific polyclonal and monoclonal antibodies. Synthetic peptides having amino acid sequence identity to 10 or more consecutive amino acid residues of a MKS1 protein can be synthesised and used as antigen for the production of specific MKS1 antibodies. It is common to couple the synthetic peptide to a carrier protein, e.g. PPD (Purified Protein Derivative; Bardarov *et al.* 1990, *FEMS Microbiology Letters* 71: 89-94), to enhance the stability of the antigen and improve the presentation of the antigen to the immune humoral response system. Polyclonal and monoclonal antibodies can be raised, screened and tested according to standard protocols, as given by Harlow and Lane (1988) *Antibodies, A Laboratory Manual*, Cold Spring Harbour Publ. NY. A variety of immunoassay formats may be used to select antibodies specifically immunoreactive for a protein. For example, solid-phase ELISA immunoassays, immunoblots, or immunohistochemistry are regularly used for this purpose. Typically a

specific or selective reaction will be at least twice background signal or noise and more typically more than 10 to 100 times background.

### Examples

- 5 MPK4 is a plant protein kinase whose regulatory functions include default repression of SA-dependent SAR, a pathway that primarily mediates resistance to certain biotrophic pathogens via PR gene expression. In addition, MPK4 is involved in the activation of PDF1.2 expression in response to jasmonate and ethylene, pathways that mediate resistance
- 10 against necrotrophs and wounding herbivores (Petersen et al., 2000 *supra*). Since the regulatory functions of MPK4 are dependent on its kinase activity, it is likely that MPK4 interacts with and phosphorylates protein substrates which directly or indirectly lead to the control of gene expression appropriate to various pathogen responses. Hence, the identification and isolation of a
- 15 protein, which interacts with and is phosphorylated by MPK4, would provide a key regulator of SA-dependent SAR in plants.

### Example 1

#### ***Arabidopsis* MAP Kinase Substrate 1 (MKS1) interacts with MPK4**

- 20 A yeast two-hybrid screen was employed to identify proteins that interact with the MPK4 protein. The yeast two-hybrid screen, first described by Fields and Song in 1989 (*Nature* 340: 245-24) is a common method used to detect protein-protein interactions. This screen exploits inherent properties of transcription factors, namely that are composed of two separate domains: a
- 25 DNA-binding domain and a transcription activation domain. A physical association of the two domains of a transcription factor is required in order for it to bind to a promoter and activate transcription of a downstream gene. DNA sequences encoding fusion proteins, comprising the DNA-binding domain or the activation domain of a transcription factor, can be constructed and co-
- 30 expressed in yeast. Interaction between the two fusion proteins will result in a functional transcription factor. If a DNA binding domain-MPK4 fusion protein



(bait), and an activation domain fused to an MSK1 interacting protein (prey) are simultaneously expressed in yeast, a functional interaction between the two fusion proteins can be detected by the transcription of nutritionally essential genes and reporter genes cloned in yeast. The yeast two-hybrid screen is commonly based on the detection of yeast colonies in which transcription of these essential genes enables cell growth on histidine- or leucine-deficient media, and detectable  $\beta$ -galactosidase activity.

An *Arabidopsis* cDNA library fused to the activation domain of a transcription factor (prey) was screened for potential MPK4 interacting partners using the following yeast two-hybrid system. *Saccharomyces cerevisiae* strain PJ69-4A (MATa trp1-901 leu2-3, 112 ura3-52 his3-200 gal4 gal80 LYS2::GAL1-HIS3 GAL2-ADE2 met2::GAL7-lacZ; (James *et al.*, 1996, *Genetics* 144: 1425-1436) was used as host strain for two hybrid screening. Cells were grown at 30°C in liquid YPD medium ([www: clontech.com](http://www.clontech.com)) or on YPD agar plates. Transformed yeast cells were grown in liquid SD medium or on SD agar (Minimal SD Agar Base; [www:clontech.com](http://www.clontech.com)) plates supplemented with drop-out supplements ([www:clontech.com](http://www.clontech.com)) lacking specific amino acids. Yeast cells were transformed using the lithium acetate/polyethylene glycol method (Ito *et al.*, 1983, *J Bacteriol* 153: 163-168). Library screening was performed with the MPK4 bait encoded by the full-length *MPK4* cDNA from *Arabidopsis thaliana* Ecotype Ler, cloned into the *Bam* H1 site in pGBD-C1 (James *et al* 1996, *supra*). Both GAL4-based library screens were performed with the *Arabidopsis* MATCHMAKER cDNA library cloned in pGAD10 GenBank #U13188 ([www: clontech.com/techinfo/manuals](http://www.clontech.com/techinfo/manuals)). Two independent screens of the library were conducted with the MPK4 bait, and in total the number of screened clones ( $6 \times 10^7$ ) covered the library 20 times. Subsequently 7.4 million colonies were screened with MKS1 as bait, corresponding to 25 times the number of individual clones in the library.

A single full-length cDNA, designated MAP Kinase Substrate 1 (MKS1), corresponding to the intron-less *Arabidopsis* gene At3g18690, was found to interact with MPK4 in the yeast two-hybrid screen, shown in Figure 1A. A similar interaction was observed after switching MPK4 and MKS1 as prey and bait, respectively. To test the specificity of the MAP kinase interaction, the interaction of MKS1 with other plant MAP kinases was tested in the yeast two-hybrid assay. The following MAP kinase cDNA sequences were cloned as AD fusions in pGAD424 (www: clontech.com): MPK3 (nucleotides 149-1261 of NM\_114433) using *Bam*HI/*Sa*I sites; MPK5 (nucleotides 466-1218 in NM\_117204) using *Nco*I/*Not*I sites; MPK6 (nucleotides 116-1303 in NM\_129941) using *Bam*HI/*Sa*I sites; MPK17 (nucleotides 1-1740 in NM\_126206) using *Nco*I/*Not*I sites. In contrast to MPK4, the MPK3, 5, 6 or 17 (Ichimura *et al.* 2002, *Trends in Plant Sci.*, 7, 301-308) preys did not interact with the MKS1 bait (Figure 1A), confirming the specificity of the MKS1-MPK4 interaction.

MKS1 is a protein of 222 amino acid residues having a predicted molecular mass of 24 kDa, and the sequence of Seq ID No: 2 (At3g18690). MKS1 is encoded by nucleotides 80 to 748 of the *Arabidopsis* gene At3g18690; GI: 18401970 (SEQ ID No: 1). MKS1 contains 11 putative MAP kinase phosphorylation sites (Ser-Pro), indicated in Figure 1B, based on sequence homology to other described phosphorylation sites (minimal consensus sequence S/TP; Sharrocks *et al.*, 2000, *Trends in Biochem Sci.*, 25: 448-453). The coding sequence for MKS1 was used in a standard protein-protein BLAST and tblastn search against the database at the www.ncbi.nlm.nih.gov/blast/BLAST.cgi and www.arabidopsis.org/Blast sites. The BLAST searches identified the following nucleic acid sequences comprising ORFs coding for previously unknown proteins, now identified as: *Arabidopsis* MKS1 gene homologue (Seq ID No: 5; Acc.No:At3g21326) encoding MKS1 protein homologue (Seq ID No: 6), *Brassica oleracea* MKS1 gene homologue (Seq ID No: 9; Acc.No:BH544707; GI:17796488) encoding

MKS1 protein homologue (Seq ID No: 10), *Brassica oleracea* (Seq ID No: 13; Acc. No:BOHBT92TR + BOGQI24TF) encoding MKS1 protein homologue (Seq ID No: 14), *Glycine max* MKS1 gene homologue (Seq ID No: 15; Acc.No: BE020960) encoding MKS1 protein homologue (Seq ID No: 16),  
 5 rice MKS1 gene homologue (Seq ID No: 19; Acc.No:CAD40925; GI: 21740554) encoding MKS1 protein homologue (Seq ID No: 20), rice MKS1 gene homologue (Acc.No: OsAP004654) encoding MKS1 protein homologue (Seq ID No: 26), maize MKS1 gene homologue (Acc.No: ZmCC613160) encoding MKS1 protein homologue (Seq ID No: 27), maize MKS1 gene  
 10 homologue (Acc.No: ZmCC635639) encoding MKS1 protein homologue (Seq ID No: 28, which all share sequence identity with *Arabidopsis* MKS1 (Seq ID.No: 2) and comprise Domains 1 and 2, as shown in the protein alignment given in Figure 1B. The alignment was generated with the aid of CLUSTAL programs (clustalw.genome.ad.jp; Jeanmougin,F. *et al.*, (1998) *Trends*  
 15 *Biochem Sci*, 23, 403-5; Thompson,J.D., *et al.* (1997) *Nucleic Acids Research*, 24:4876-4882; Higgins, D. G., *et al.* (1996) Using CLUSTAL for multiple sequence alignments. *Methods Enzymol.*, 266, 383-402.  
 5) Thompson, J.D., Higgins, D.G. and Gibson, T.J. (1994) *Nucleic Acids Research*, 22:4673-4680; Higgins,D.G., *et al.* (1992) *CABIOS* 8,189-191;  
 20 Higgins,D.G. and Sharp,P.M. (1989) *CABIOS* 5,151-153; Higgins,D.G. and Sharp,P.M. (1988) *Gene* 73,237-244). Furthermore, MKS1 homologues or orthologues are also found in maize (*Zea mays*), tobacco (*Nicotiana tabacum*) and clover (*Medicago truncatula*) (Figure 1B). *Arabidopsis* MKS1 protein (At3g18690) shares a sequence identity of 84.8% with *Brassica oleracea* MKS1 (Acc.No:BH544707) and 78.4% with *Brassica oleracea* (Acc no: BOHBT92TR + BOGQI24TF). The identified MKS1 homologues all comprise amino acid sequence domains 1 and 2, or conservatively modified variants thereof.

## 30 Example 2

***Arabidopsis* MPK4 interacts with and phosphorylates MKS1 *in vitro***

**A. MPK4-MKS1 interaction *in vitro***

To substantiate the interaction between MPK4 and MKS1, detected in the yeast two-hybrid screen, *in vitro* interaction assays (pull-down assays) were performed with recombinant MPK4 and MKS1 proteins. Recombinant MKS1 was obtained by bacterial expression according to the following procedure. The full-length MKS1 coding sequence (At3g18690 nucleotides 80 to 748) was cloned in-frame with the glutathione-S-transferase (GST) gene in the *Xho* I site of pGEX-5X plasmid ([www: amershambiosciences.com](http://www.amershambiosciences.com)). Expression of the recombinant protein in *E. coli* BL21 (pLysS) cells (www:novagen.com) was induced with 0.1 mM isopropyl- $\beta$ -D-thiogalactopyranoside (IPTG) at 30°C for 3–4 h, and 2% ethanol was added before induction. GST protein was similarly expressed in *E. coli* from the pGEX-5X plasmid. GST and GST-fusion proteins were purified from whole cell extracts of *E. coli* by binding to glutathione–Sepharose 4B beads (www: amershambiosciences.com), in the presence of proteinase inhibitors (2  $\mu$ g/ml leupeptin, 1 mM AEBSF (4-(2-Aminoethyl)-benzenesulfonylfluoride.HCl), 2  $\mu$ g/ml antipain, 5 mM EDTA, 5 mM EGTA, 2  $\mu$ g/ml aprotinin). Proteins used in pull-down assays were not eluted from the glutathione sepharose beads. <sup>35</sup>S-methionine-labelled MPK4 was generated by coupled transcription-translation of the bait plasmid pGBKT7-MPK4 from the two hybrid screen, using a T7 coupled reticulocyte lysate system ([www: promega.com/tbs/tb126/tb126.pdf](http://www.promega.com/tbs/tb126/tb126.pdf)).

Pull-down assays were performed as follows: 10  $\mu$ l <sup>35</sup>S-MPK4 was mixed with 200  $\mu$ l 1% BSA in Bead Binding (BB) Buffer (BB Buffer; 50 mM KPO<sub>4</sub> pH 7.5, 150 mM KCl, 1 mM MgCl<sub>2</sub>, 2  $\mu$ g/ml leupeptin, 1 mM AEBSF, 2  $\mu$ g/ml antipain, 2  $\mu$ g/ml aprotinin), incubated on ice for 15 min, and then centrifuged for 10 min at 4°C. The supernatant was added to 2-5  $\mu$ g GST or GST-fusion protein bound to sepharose beads in 200  $\mu$ l 1% BSA in BB Buffer and incubated for 2 hrs at 4°C with rotation. The beads were washed 3 times with 1 ml wash buffer (50 mM KPO<sub>4</sub>, pH 7.5, 150 mM KCl, 1 mM MgCl<sub>2</sub>, 10%

glycerol, 5% Triton X-100) with proteinase inhibitors and were then subjected to SDS-PAGE (Sodium dodecyl sulphate polyacrylamide gel electrophoresis; Laemmli, 1970 *Nature* 227: 680) separation on 15% gels.

- 5     The pull-down assay demonstrated that *in vitro* synthesized MPK4 (Fig. 2A, lane 1) interacts with and was bound by recombinant MKS1-GST (Lane 3), but not by GST alone (Lane 2), thereby confirming the MPK4-MSK1 interaction detected in the yeast two-hybrid screen.

10    **B. MPK4 phosphorylation of MKS1 *in vitro***

- The ability of MPK4 to phosphorylate putative MAP kinase Ser-Pro phosphorylation sites in MKS1 was investigated by *in vitro* phosphorylation assays. Full-length and C-terminally truncated histidine-tagged MSK1 were expressed and purified from *E. coli*. MKS1 nucleotide sequence (nucleotides  
15   80 to 748 of At3g18690) encoding full-length MKS1 protein, was cloned into the *Xho* I site of the pET15b plasmid ([www:novagen.com](http://www.novagen.com)). Nucleotide sequences encoding MKS-1 with terminal deletions, C1-C3, were constructed by restriction digest of the MKS1-HIS containing pET15b vector. The C1 deletion was generated with *Bst*BI and *Bpu*1102I, the C2 deletion  
20   with *Nhe*I and *Bpu*1102I, and the C3 deletion with *Sty*I and *Bpu*1102. The digested plasmids were end-filled by incubation with 3 U Klenow enzyme and 10  $\mu$ M deoxyribonucleotides for 30 minutes at 37°C, and then re-ligated by overnight incubation with ligase enzyme at 16 °C. The plasmid constructs encoding full-length MKS1 (amino acids 1-222), C1-MKS1 truncation (amino  
25   acids 1-196), C2-MKS1 truncation (amino acids 1-123) and C3-MKS1 truncation (amino acids 1-73), as shown in Figure 1B, were transformed into *E. coli* BL21 (pLysS) cells ([www: novagen.com](http://www.novagen.com)), expression was induced with 0.1 mM isopropyl- $\beta$ -D-thiogalactopyranoside (IPTG) at 30°C for 3–4 h, adding 2% ethanol prior to induction. Expressed MKS1 was extracted using  
30   BugBuster and Benzonase assisted protein extraction, and purified by affinity

binding of the histidine tag to Ni-NTA resin, according to the instructions of the manufacturer (www.novagen.com).

HA (influenza hemagglutinin antigen)-tagged MPK4 (HA-tag is 6 X YPYDVPDYA) was expressed by transgenic *Arabidopsis* plants (Petersen *et al.*, 2000, *supra*). The HA-MPK4 was purified from protein extracts of the plants as described (Romeis *et al.*, 1999, *Plant Cell* 11: 273-287), except that a buffer change was not made prior to immunoprecipitation (Romeis *et al.*, 1999, *Plant Cell* 11: 273-287). 100 µg of total protein was immunoprecipitated from the plant protein extract with 2 µg/ml monoclonal 12CA5 HA-antibody (Boehringer) by affinity to the HA tag. Protein concentrations were determined with the Bradford dye-binding procedure (Bradford, 1976, *Anal Biochem* 131: 248-254). The resulting sepharose beads, with immunoprecipitated MPK4, were washed in kinase buffer (200 µM ATP, 80 mM Tris-HCl, pH 7.5, 8 mM EGTA, 120 mM MgCl<sub>2</sub>, 4 mM Na<sub>3</sub>VO<sub>4</sub>, 4 mM DTT) to remove the immunoprecipitation buffer and suspended, as a 50% slurry, in kinase buffer.

Phosphorylation assays were performed by mixing 10 µl MPK4-sepharose slurry, 5 µg substrate protein and 0.4 µl 300 µM <sup>32</sup>P-γ-ATP (3 µCi) with kinase buffer in a final volume of 30 µl. The assay samples were incubated for 1 h with agitation at 30°C, where after the assay proteins were separated by SDS-PAGE, and the gels subsequently dried on Whatmann 3MM paper and the radiolabelled proteins detected on a phosphorimager screen.

HA-tagged MPK4, immunoprecipitated from *Arabidopsis* plants, is shown to *in vitro* phosphorylate MKS1 as efficiently as myelin basic protein (MBP; Sigmasource), which is a standard MAP kinase substrate, as shown in Figure 2B, lanes 1 versus 5). Immunoprecipitated extracts of non-transgenic *Arabidopsis* plants (wt) failed to phosphorylate MKS1 (Figure 2B, lanes 6-8) confirming that the HA-antibody specifically immunoprecipitates HA-tagged MPK4. Furthermore, a mutant HA-tagged MPK4, with substitutions in the

kinase activation loop abolishing MPK4 activity (T201A/Y203F; Petersen *et al.*, 2000, *supra*) was similarly found not to phosphorylate MKS1 or MBP.

In order to identify which sites in MKS1 are phosphorylated by MPK4, C-terminal MKS1 truncations (C1, C2, C3), lacking some of the putative Ser-Pro phosphorylation sites (Figure 1B), were tested in the phosphorylation assay. HA-tagged MPK4 readily phosphorylated both full-length and C-terminal MKS1 truncations, including C3 MKS1, which retains only 2 putative phosphorylation sites (Ser30 and Ser72), as seen in Figure 2B, lanes 2-4. In order to map the functional phosphorylation sites in the C3 MKS1 protein, the encoded MSK1 sequence was altered from Ser30 to Ala30 (S30A) by *in vitro* mutagenesis, by substituting the codon TCA for GCA in the full-length and C3 truncated *MKS1* gene. Although the mutant C3 truncated MKS1 (C3-S30A) was not phosphorylated by HA-tagged MPK4, the mutant full-length MKS1 (S30A) was phosphorylated (Figure 2C, lanes 1 and 2 versus 3 and 4). This indicates that MPK4 phosphorylates MKS1 at Ser30, as well as other additional sites in the MKS1 protein.

A synthetic 22 amino acid peptide (Pep22), corresponding to amino acid residues 13-35 of MKS1 and comprising Ser30, shown as in Figure 1B, was synthesized by KJ Ross ([www.tagc.com](http://www.tagc.com)). Pep22 is an efficient competitor of full-length MKS1 for phosphorylation by MPK4, when added to the *in vitro* assay in a molar ratio of 1:1 (Pep22:MKS1) as shown in Figure 2D, top. The Flg22 peptide, with amino acid sequence QRLSTGSRINSAKDDAAGLQIA, which is known to activate immediate pathogen responses via the flagellin receptor, involving MPK3, 5, 6 and 17 as well as WRKT 22 and 29 (Asai *et al.* 2002, *Nature* 415: 977-983), was used as a control in this assay. Since the Flg 22 peptide did not compete MKS1 phosphorylation (Figure 2D, bottom) it is likely that the Pep22 domain of MKS1 specifically interacts with MPK4.

### C. MPK4 interacts with the N-terminal region of MKS1

The location and sequence specificity of the interaction between MPK4 and MKS1 was investigated employing the yeast 2-hybrid system described in Example 1. Nucleic acid molecules encoding the three C-terminal truncated forms of MKS1, (see Example 2B) as well as an N-terminal truncated form, N1-MKS1 (amino acids 55-222; deleted region is indicated as N1- in Table 1) were cloned into the 2-hybrid vectors and screened as MPK4 interacting partners. Only the N-terminally deleted MKS1 failed to interact with MPK4, confirming that the N-terminal 54 amino acid region is essential for MPK4-MKS1 interaction. N-terminal amino acid residues in MKS1 that are essential for this interaction were examined by testing the ability of MPK4 to interact in the 2-hybrid system with mutant MKS1 polypeptides. Site-directed mutagenesis was used to generate nucleic acid molecules encoding 18 full-length mutant MKS1 polypeptides each having a single amino acid substituted with alanine (denoted 'A' in Table 1), localised in the N-terminal domain 1 and pep22 regions.

### Table 1

```

      Pep22                               N1-A                               Δ-C3
MKS1: MDPSEYFAGGNPSDQONQKRQLQICGPRPSPFSVHKDSHKIKKPPKHPAPPPNRPDPFPYIPREPVVIIYAVSPKVVHATASEFMNVVQRLTG
I mutants*:      A      AAA A A AA A AA AA
N-I mutants*:      A      AA AA
                        i qrrp pl v dsh ikk                                pviiv SPkvvha eFm vVOrLTG
                        DOMAIN I                                              DOMAIN II

```

20

Mutation of five amino acid residues in the N-terminal region of MKS1, corresponding to the pep22/domain 1 region was found to prevent the interaction between MKS1 and its kinase MPK4. This domain I, essential for MPK4 interaction, shares amino acid sequence homology with MAPK docking sites (Sharrocks *et al.*, 2000, Trends Biochem. Sci. 25: 448-453).



### Example 3

#### Antibodies for MKS1 detection

To provide tools for the detection of MKS1 expression *in vitro* or *in vivo* in single or multicellular organisms, polyclonal (pa-Pep22) and monoclonal antibodies (ma-Pep22 & ma-Pep22p) were raised against the peptide Pep22 (SDQQNQKRQLQICGPRPSPLSVH), corresponding to amino acid residues 13-35 of MKS1. Ten to twelve-week old female Balb/cCF1 F1-hybrid mice were used to raise both polyclonal and monoclonal antibodies. The mice were primed with 0.2 mL live BCG vaccine, delivered intraperitoneally. One month later the mice were immunised with the antigen Pep22 coupled to PPD (Purified Protein Derivative; Bardarov *et al.* 1990, *FEMS Microbiology Letters* 71: 89-94), absorbed onto the adjuvant Al(OH)<sub>3</sub>. The total volume of vaccine per immunisation was 500 µL, containing 15 µg of PPD and 1 mg of adjuvant. The antigen was injected intraperitoneally at 2-week intervals. To prepare polyclonal antibodies from the immunised mice, blood samples were collected 10 days after each immunisation and assayed for specific recognition of HIS-tagged MKS1 protein, expressed and purified from *E. coli*, followed by SDS-PAGE separation and semi-dry transfer and immunoblotting (Current protocols, [www:wiley.com](http://www.wiley.com)). Western blots were developed using alkaline phosphatase conjugated anti-mouse antibody (Promega). Monoclonal antibodies were prepared from immunised mice found to produce positive antisera, essentially as described by Kohler and Milstein (1975) in *Nature* 256: 495-497, as modified by Reading (1982) in *J Immunol Methods* 53: 261-291. After hybridoma cell fusions, culture supernatants were tested for specific recognition of HIS-tagged MKS1 protein, by enzyme-linked immunoabsorbent assay (ELISA; Current protocols, ([www: wiley.com](http://www:wiley.com))) and immunoblotting as described above for polyclonal antibodies.

Polyclonal antibody, pa-Pep22, specifically recognised MKS1 present in extracts of *E. coli* and wild type *Arabidopsis* plants, as shown by Western blotting in Figure 3A. The same result was obtained using the monoclonal

antibody ma-Pep22 (not shown). Monoclonal antibody ma-Pep22 (HYB 330-01), specifically recognised and immunoprecipitated MKS1 present in extracts of wild type *Arabidopsis* plants, since the immunoprecipitated MKS1 was detected by the pa-Pep22 polyclonal antibody, as seen in lane 1 (upper band) of a Western blot (Figure 3B). The lower band is due to binding of the secondary anti-IgG antibody to the ma-Pep22 light chain, which was also present in a control immunoprecipitation with ma-Pep22 where plant extract is omitted, seen in lane 2 (Figure 3B).

#### 10 **Example 4**

##### ***Arabidopsis* MPK4 interacts with MKS1 *in vivo***

Interaction between MKS1 and MPK4 *in vivo* in *Arabidopsis* plants was demonstrated by the ability of the MKS1 specific monoclonal antibodies to co-immunoprecipitate MPK4 with MKS1 from leaf extracts. Leaf protein extracts were prepared as described in Example 2B, from transgenic *mpk4* plants complemented to wild type by a functional HA-tagged MPK4 gene (Fig. 3C; Petersen *et al.*, 2000, *supra*). Immunoprecipitates of the leaf extracts were analysed by SDS-PAGE and Western blots, which were probed with anti-HA antibody to detect HA-tagged MPK4. As shown in Figure 3C lane 1, ma-Pep22 monoclonal antibody co-immunoprecipitated HA-tagged MPK4, which was detected with the anti-HA antibody. The co-immunoprecipitated HA-tagged MPK4 co-migrated with MPK4 immunodetected in whole plant extracts (lane 1 versus lane 3). Monoclonal antibody (ma-Con), that does not detect MKS1 in plant extracts, was unable to immunoprecipitate MPK4 (lane 2). The upper bands immunodetected in lanes 1, 2 and 4 of Figure 3C are likely due to binding of the secondary anti-IgG to the heavy chain of the immunoprecipitating monoclonal IgGs. Only the upper band was detected in a mock-plant extract containing MaPep22 (lane 4).

The *in vivo* phosphorylation status of MKS1 in extracts wild-type *Arabidopsis* leaves (*Ler*) as compared with *mpk4* leaves, was examined following immunoprecipitation of MKS1 with ma-Pep22 monoclonal antibody. The MKS1 polyclonal antibody ( $\alpha$ -MKS1) detected equal amounts of MKS1  
5 immunoprecipitated from wild-type and *mpk4* plants, as shown in figure 3, D. However, a phosphoserine/phosphothreonine-specific antibody ( $\alpha$ -pS/TP) only detected MKS1 in wild-type plants, thereby confirming that MKS1 is a substrate for MPK4 and is a key component of the signal transduction pathway in plants which facilitates a disease response.

10

### Example 5

#### Transgenic *Arabidopsis* plants with modified MKS1 expression

Transgenic plants, expressing elevated or reduced levels of MKS1 protein, were generated in *Arabidopsis thaliana* ecotype Landsberg erecta (*Ler*) via  
15 *Agrobacterium*-mediated transformation, according to the floral dip method (Clough and Bent, 1998 *Plant J.*, 16:735-43). Transgenes were inserted between the T-DNA borders of pCAMBIA binary vectors, comprising the NPTII (kanamycin) resistance gene, and then transformed into *Agrobacterium*, and stably integrated into the *Arabidopsis* genome.

20

Constitutive over-expression of MKS1 in plants was obtained by the stable integration of CaMV 35S-MKS1 transgenes in *Arabidopsis*. The *Arabidopsis* MKS1 coding sequence (nucleotides 80-748 of Seq ID. No: 1 (At3g18690)) was amplified from its respective gene by PCR using a 5' primer (Seq ID.No:  
25 3) and 3' primer (Seq ID No: 4). A transgene comprising the CaMV 35S promoter sequence (GI: 2173396; with Seq ID. No: 23), operably linked to a MKS1 coding sequence (nucleotides 80-748 of At3g18690), was generated by replacing the GUS ORF in pCAMBIA1301 (AF234297) by the MKS1 sequence, ligated with *Nco* I/*Bst* EII linkers. *Arabidopsis* transformants were  
30 selected by resistance to the antibiotic hygromycin incorporated into the seedling growth medium. Transformants with an integrated copy of the CaMV

35S-MKS1 transgene in the *Arabidopsis* genome were identified by northern blotting with a MKS1 probe (nucleotides 80-748 of At3g18690) and western blotting with maPep22.

- 5 Self-fertilisation of *Arabidopsis* transformants, with an integrated copy of the CaMV 35S-MKS1 transgene, led to seed formation, with the stable inheritance of the transgene in the progeny, and subsequent generations. Cross-pollination of the primary transformed plants or their progeny with control, non-transformed plants, generated progeny that inherited the
- 10 transgene according to Mendelian genetics.

Silencing of MKS1 expression in plants by RNA interference (Chuang and Meyerowitz, 2000, *Proc Natl Acad Sci. U S A.* 97: 4985-4990) was obtained by the stable integration of a CaMV 35S-MKS1 RNAi transgene. The MKS1

- 15 coding sequence (nucleotides 80-748 of Seq ID No:1 (At3g18690)) was first inserted in the plasmid SLJ1382B1 (Andrea Ludwig and Jonathan DG Jones, Sainsbury Laboratory, UK), derived from plasmid SLJ4D4 (Jones *et al.* 1992, *Transgenic Research* 1: 285-297). The MKS1 coding sequence was cloned, in opposite orientations, on either side of an intervening intron in
- 20 pSLJ1382B1, as 5' *Xba* I/*Fse* I- and 3' *Asc* I/*Xho*-linked fragments. The intron had the sequence:

GTAAGTTTCTGCTTCTACCTTTGATATATATAATAATTATCATTAATTAG  
TAGTAATATAATATTTCAAATATTTTTTTCAAATAAAAGAATGTAGTATAT  
AGCAATTGCTTTTCTGTAGTTTATAAGTGTGTATATTTTAATTTATAACTTT  
25 TCTAATATATGACCAAAATTTGTTGATGTGCAG and Seq ID. No: 25. The resultant RNAi cassette was excised with *Eco* RI and *Hind* III and cloned into corresponding sites in pCAMBIA3300 (derivative of pCAMBIA1201, AF234293). This construct was transformed into ecotype Col-0, and transformants were selected by resistance to spraying with the herbicide

- 30 BASTA (Glufosinate-ammonium; [www: bayercropscience.com](http://www.bayercropscience.com)).

Transgenic plants with elevated levels of MKS1, expressed under control of the constitutive promoter CaMV 35S (35S-MKS1), were identified by immunodetection of MKS1 in plant protein extracts analysed by western blotting with the polyclonal antibody pa-Pep22, as shown in Figure 4A.

- 5 Transgenic plants in which MKS1 expression was silenced by RNA interference were similarly identified by immunodetection of MKS1 levels in plant protein extracts (Figure 4A). The 35S-MKS1 transgenic plants exhibited semi-dwarfism in contrast to the dwarf habit of *mpk4* mutants (Figure 4B). The MKS1-RNAi plants were phenotypically wild type in their growth habit  
10 (not shown).

#### Example 6

##### Properties of transgenic *Arabidopsis* plants with modified MKS1 expression

- 15 **A. MKS1 regulates the expression of pathogen resistance genes, but not wound and methyl jasmonate response genes, in plants.**

The steady-state levels of MKS1, PR and wound-induced gene transcripts were measured in total RNA samples extracted from *Arabidopsis* plants  
20 which were analysed by Northern blotting and hybridization with DNA probes according to standard protocols. DNA probes were amplified by PCR, with sequence-specific primers, from the following cDNA or genomic DNA templates: MKS1 (nucleotides 80-748 of At3g18690), PR1 (nucleotides 84-530 in M90508), *PDF1.2* (EST 37F10T7), *VSP* (nucleotides 3-236 in  
25 ATTS0751/GBGA288), WR3 (WR3 probe in AtT5G50200, described by Leon J. *et al.*, 1998, *Mol Gen Gen* 258: 412-419).

35S-MKS1 transgenic plants accumulated elevated levels of MKS1 mRNA compared to wild-type, consistent with increased MKS1 synthesis in these  
30 plants (Figure 5A). Levels of the pathogen resistance PR1 mRNA, were

enhanced in 35S-MKS1 transgenic plants and in *mpk4* mutant plants when compared to wild-type plants (Figure 5A, lane1 versus 2 and 3).

Comparative transcript profiling of wild type, *mpk4*, and 35S-MKS1 plants  
5 was used to examine differences in global gene expression with the  
Affymetrix ATH1 array covering 22,810 transcripts. Total RNA was isolated  
from three replicates of wild type *Ler*, *mpk4*, and 35S-MKS1 grown in an  
chamber with 16hr light (21C) and 8hr dark (16C). RNA was amplified  
according to the standard Affymetrix protocol and hybridized to the Affymetrix  
10 ATH1 oligonucleotide microarray (Acc. # E-MEXP-173, ArrayExpress  
database, EBI). Raw intensity data was normalized using R implementation  
of q spline (Gautier et al., 2004, Bioinformatics 20: 307-315). An  
implementation in the statistical language R of the logit-t method (Lemon et  
al., 2003, Genome Biol. 4: R67) applying one-way ANOVA was used to  
15 calculate statistical significance for differentially gene expression (R source  
code available on request). Genes with p-value less than 0.01 were  
considered significant. Fold change =  $\max(\text{over all } j (\text{median}(E_{ij})) - \min(\text{over all } j (\text{median}(E_{ij})))$ , where  $E_{ij}$  is the  $ij$ 'th gene expression index value,  $j$  is the  
genotype, and  $i$  the sample. Gene expression index values were calculated  
20 using the PM only implementation method (Li & Wong, 2001, Genome Biol.  
2: R0032.1-11). Gene expression profiles for the 800 most significantly  
differentially expressed genes were clustered by k-means. Examples of over  
expressed genes in 35S-MKS1 are shown in Table 2.  
In the 35S-MKS1 lines *MKS1* was the most significantly differentially  
25 expressed gene due to its over-expression. Significantly, mRNAs of  
pathogen responsive genes including *PR1*, *PR2* and the SA biosynthesis  
enzyme ICS1 were increased in both *mpk4* and 35S-MKS1 plants as  
compared to wild-type.

Table 2

	Fold	Accession	Annotation		Fold	Accession	Annotation
	59.7	At3g18690	MKS1		29.7	At4g01370	MAP kinase 4 (MPK4)
	34.1	At5g39100	Germin-like protein (GLP6)		21.2	At1g53870	Hypothetical protein
5	13.2	At2g43570	Endochitinase isolog		20.6	At5g44420	antifungal protein (PDF1.2)
	11.1	At2g14610	PR1-like protein		5.4	At3g25760	Hypothetical protein
	9.5	At3g22600	Lipid transfer protein		4.3	At2g36145	Expressed protein
	8.1	At1g75040	Thaumatococcus-like protein (PR5)		4.1	At1g20190	Expansin S2 precursor
	6.1	At1g74710	Isochorismate synthase (ICS1)		3.9	At1g21500	Unknown protein
	5.3	At1g69930	Glutathione transferase		3.8	At4g26530	Fructose biphosphate aldolase
10	4.2	At3g57260	Beta-1,3-glucanase (PR2)		3.6	At1g04800	unknown protein

MKS1, in contrast to MPK4, is shown not to be involved in the response to wounding and necrotrophic attack in plants. Plants respond to wounding and necrotrophic attack by the transcriptional activation of jasmonate and/or ethylene responsive genes including *VSP*, *WR3* and *PDF1.2*, in which MPK4 is known to play a regulatory role (Petersen *et al.*, 2000, *supra*; Andreasson E. and Mundy J, unpublished). The steady-state levels of these wound-induced genes was determined in *Arabidopsis* plants, subjected to wounding by making 1 to 3 cuts over the mid vein with a pair of scissors. *VSP* and *WR3* mRNAs were induced in wild-type plants within 2 hours of wounding (Figure 5B, lane 7), but were undetectable or greatly reduced in *mpk4/ NahG* plants expressing the bacterial salicylate hydroxylase that degrades SA (Figure 5B, lane 3). The same results were also seen following wounding of the *mpk4* mutant (not shown). Silencing MKS1 expression in RNAi-MKS1 plants did not prevent a wild-type wounding response with the accumulation *VSP* mRNA (Figure 5C, lanes 2 versus 4). These results indicate that MKS1 is not required for wound-responsive *VSP* expression. Silencing or over-expression of MKS1 did not significantly affect the levels of PDF2.1 mRNA accumulation following 48hr of MeJA treatment (Fig. 5D). This indicates that MKS1, in contrast to MPK4 (Petersen *et al.* 2000 *supra*) is not required for MeJA responsive PDF1.2 expression.

**B. Salicylic acid levels are enhanced in 35S-MKS1 transgenic plants**

The steady-state free and glycosylated salicylic acid content of *Arabidopsis* plants was analysed in plant extracts prepared by grinding plant tissue in liquid nitrogen, extracting the ground tissue in methanol, following by an  
5 ethylacetate:cyclopentane:isopropanol partition of the extract according to (Newman *et al.*, 2001, *Mol Plant-Microbe Interactions* 14: 785-792). The salicylic acid content was analysed by HPLC using a diode array detector between 180-350nm, as previously described Newman *et al.*, 2001, *supra*. Salicylic acid levels were significantly elevated in 35S-MKS1 transgenic  
10 plants in comparison to wild-type plants, as shown in Figure 6A.

**C. Pathogen resistance is enhanced in 35S-MKS1 transgenic plants**

Resistance to the plant pathogen *Pseudomonas syringae* is shown to be  
15 controlled by MKS1 expression levels in transgenic plants. Four-week-old *Arabidopsis* plants were infiltrated with a suspension of  $1 \times 10^5$  cfu/ml of virulent *Pseudomonas syringae* pv. *tomato* DC 3000 strain. Bacterial growth on infected plants was subsequently assayed by grinding four 0.5 cm<sup>2</sup> leaf pieces in 10mM MgCl<sub>2</sub> for each sample. Dilutions were distributed on NYG  
20 agar plates containing rifampicin, cycloheximin and kanamycin, and colonies were counted, as previously described (Parker *et al.* 1996, *Plant Cell* 8: 2033-2046). 35S-MKS1 transgenic plants exhibited increased resistance to *P. syringae* DC3000, as seen for *mpk4* plants, as shown in Figure 6B. The disease response of 35S-MKS1 transgenic lines expressing different levels of  
25 MKS1, indicated that MKS1 expression is directly correlated with PR1 expression and resistance to *Pseudomonas* attack (data not shown). In contrast, of MKS1-RNAi plants were significantly less resistant to *P. syringae* DC3000 than wild type plants (Figure 6C) confirming the key role of MKS1 in the development of SAR.



### E. Localisation of MKS1 expressed in transgenic plants

Green fluorescent protein (GFP) expressed in plant cells can be detected by virtue of its fluorescent properties, and GFP-protein fusions have provided a valuable tool for determining the whole plant and subcellular expression pattern of proteins of interest (Stewart, 2001, *Plant Cell Rep.* 20:376-82). *Arabidopsis* plants transformed with MKS1-GFP gene fusions, under control of CaMV 35S or MKS1 promoters, were generated to determine MKS1 cellular localisation. The MKS1 coding sequence with *Eco* RI linkers (nucleotides 80 to 748 in At3g18690) was N-terminally fused in frame with a GFP coding sequence, operably linked to a CaMV 35S promoter in the binary vector pCAMBIA 1302 (AF234297). The MKS1-GFP gene fusion cloned in pCAMBIA 1302 was placed under the control of the MKS1 promoter by substituting the CaMV 35S promoter by a 1.9 kb MKS1 promoter fragment (complement of nt 15531-13589 of BAC MVE11) having *Nco* I/*Bst* EII linkers. The MPK4-GFP fusion was made by cloning a *Not*I linker genomic fragment including 1150 bp promoter region from Ler genomic DNA cloned into pAVA393 (Arnim *et al.*, 1998 *Gene* 221: 35-45). The control 35S-GUS-GFP fusion was included in the pCAMBIA1302, a derivative pCAMBIA1303. The transgenes in the binary vectors were transformed into *Arabidopsis* by *Agrobacterium*-mediated transformation and transgenic lines were selected as described in Example 5.

GFP expressed in mesophyll cells of young leaves of the transformed lines was visualised by confocal microscopy. GFP fluorescence was detected with a Zeiss LSM 510 laser-scanning microscope applying the 488 nm line of the argon laser and the corresponding dichroic mirror and a 505-530 nm band-pass filter. The generated images of GFP fluorescence in cells are vertical projections of variable numbers of optical sections.

The phenotype of the transgenic lines expressing the MKS1-GFP fusion protein was similar to that of the 35S-MKS1 transgenic lines expressing

enhanced MKS1 levels. This indicates that MKS1 retains functional activity when expressed as a GFP-MKS1 fusion protein. Similarly, the MPK4-GFP fusion protein is functional and correctly targeted when expressed in transgenic plants, since it is able to complement the *mpk4* mutant to wild type (Brodersen, Mattsson and Mundy, unpublished data). The GUS-GFP fusion protein, which lacks any specific subcellular, or extracellular targeting signals, was primarily localized to the cytoplasm of 35S GUS-GFP transgenic plants, as shown in Figure 6D. GFP-MKS1, as well as GFP-MPK4, were localised in the nucleus of mesophyll cells, consistent with the demonstrated *in vivo* interaction of these two proteins and their transcriptional repression of downstream SAR effector genes under normal growth conditions (Figure 6D).

In conclusion, transgenic plants with elevated levels of MKS1 expression show increased salicylic acid (SA) levels, PR gene expression and pathogen resistance, demonstrating that MKS1 is a key component of the SAR signal transduction pathway in plants controlling SAR and plant pathogen resistance. During negative regulation of SAR by MPK4 in wild-type plants, MKS1 is presumably phosphorylated, at one or more sites.

#### **E. MKS1 acts down-stream of MPK4 in the SAR signal transduction pathway**

MKS1-RNAi Arabidopsis plants, in which MKS1 expression is down-regulated, were crossed with *mpk4* mutants to provide double mutants with the genotype MKS1-RNAi/*mpk4*. The *mpk4* phenotype was partially suppressed in the double mutants, whereby the dwarf phenotype was reduced (Figure 7,a), the level of PR gene induction was reduced (Figure 7b) and the plants were less resistant to pathogen attack (Figure 7c). The ability of MKS1-RNAi genotype to suppress the *mpk4* phenotype confirms the essential role of MKS1, downstream of MPK4, in the SAR signal transduction pathway.

**Example 7*****Arabidopsis* MKS1 interacts with WRKY 25 and 33 transcription factors**

MKS1 is shown to be a key component of the SAR signal transduction pathway in plants, whose overexpression enhances SA levels and PR gene

5 expression. The regulatory role of MKS1 is likely to be mediated by interaction with additional down-stream members of the pathway, including transcription factors. A yeast two-hybrid screen, with MKS1-BD as the bait, was used to identify proteins capable of interaction with MKS1. The MKS1-BD fusion was constructed by inserting the full-length MKS1 coding  
10 sequence (nucleotides 80-748 of At3g18690) into the *Nco* I restriction site of pGBKT7, and transformed into *S. cerevisiae* strain PJ69-4A (www: clontech.com). A GAL-4 based library screen in yeast of *Arabidopsis* MATCHMAKER cDNA libraries was performed as described in Example 1. 7. 4 million colonies were screened with the MKS1 bait, corresponding to 25  
15 times the number of individual clones in the library. Two MKS1 interactors, the transcription factors WRKY25 (GI:15991725) and WRKY33 (GI:21105638), were identified in this screen, as shown in Figure 1A. WRKY 33 and 25 are among the 70, or more, WRKY transcription factors predicted in *Arabidopsis*, which show amino acid sequence similarity and both belong  
20 to the group I WRKYs (Eulgem *et al* 2000 *supra*). Five different truncated WRKY33 proteins interacted with MKS1 in the yeast library, the shortest corresponding to the C-terminal 188 amino acids of WRKY33. This region comprises the C-terminal WRKY domain and a region denoted the A-motif (Eulgem *et al.*, 2000, *Trends in Plant Sci* 5: 199-206).

25

The specificity of MKS1 interaction with WRKY transcription factors was examined in directed yeast two-hybrid assays with WRKY26 and WRKY29. Full-length cDNA WRKY26 (AF224699, nucleotides 23-949) and WRKY29 (AF442394, nucleotides 1-915) was fused with the nucleotide sequence  
30 encoding an AD domain in pGADT7 (www: clontech.com) using *Bam* H1 sites. WRKY26, of unknown function, is the next closest homolog to

WRKY25 and 33, while the less similar WRKY29 positively regulates innate immunity responses involving MPK3 and 6 (Asai *et al.* 2002, *Nature* 415: 977-983). However, neither WRKY26 nor WRKY29 interacted with MKS1 in this assay (Figure 1A), indicating that the interaction of MKS1 with WRKY25  
5 and 33 is specific. No activity of the reporter His3, Ade2 or lacZ gene products was detected when any fusion protein construct was co-transformed with the corresponding empty vectors (data not shown).

10 The domain in the MKS1 polypeptide required for interaction with W33 was analysed in the yeast 2-hybrid system employing vector constructs expressing the N1-MKS1, C1-MKS1, C2-MKS1 and C3-MKS1 truncated polypeptides. Since on the C3-MKS1 failed to interact with W33 in yeast this localised the interaction domain between MKS1 and W33 to the conserved domain II.

15 MKS1 is shown to be a positive regulator in the SAR signal transduction pathway, interacting with the MAP kinase MPK4 and the transcription factors WRKY25 and WRKY33. The defence response pathways, triggered by pathogen attack or wounding, which involve a series of signalling steps controlled by regulator proteins leading to the expression of resistance  
20 genes, are outlined in a model presented in Figure 7. It is proposed that the negative regulator MPK4 represses SAR by phosphorylating and interacting with MKS1 to form a complex. The MPK4-MKS1 complex may, in turn, phosphorylate the transcription factors WRKY25 and 33 that may repress transcription of a salicylic acid promoter factor. The interaction of WRKY  
25 factors with promoters (W box motifs) is known to be phosphorylation dependent (Eulgem *et al.* 2000, *supra*).

30 MKS1 is a key regulatory protein of plant SAR and thereby controls the ability of plants to survive pathogen attack. Transgenic plants expressing enhanced levels of MKS1 protein show a significantly increased level of disease resistance. Thus transgenic plants comprising a transgene expressing

enhanced levels of MKS1 may, by virtue of their increased disease resistance, produce a crop with a larger yield. Furthermore, the crop yield of these transgenic plants will be less dependent on the application of fungicides and bactericides, which are expensive and often have a negative environmental impact. The SAR response is common to many members of the plant kingdom and hence the use of MKS1 proteins to up-regulate the pathogen defence response in a wide range of plants lies within the scope of the present invention.